

**List of Lepidoptera names described as new in the scientific publications of
Márton Hreblay with information on their primary types**

ZS. BÁLINT & G. KATONA

*Department of Zoology, Hungarian Natural History Museum,
H-1088 Budapest, Baross utca 13, Hungary.
E-mails: balint@nhmus.hu, katona@nhmus.hu*

Abstract – A complete list of the 604 species-group and 28 genus-group names proposed for Lepidoptera by Márton Hreblay (1963–2000) is presented with the following data of their primary types: original description, country and published locality, sex, and depository. The genus-group name *Transeuplexia* Hreblay et Plante, 1995 is not available for any nomenclatural purpose. A short discussion about the primary type material according to their origins and depositories is given. With 7 figures and 2 tables.

Key words – list of taxon names, Márton Hreblay, Nepal, Noctuidae

INTRODUCTION

The Lepidoptera collection of Dr Márton Hreblay (1963–2000) (Fig. 1) was deposited in the Hungarian Natural History Museum by the will of his heirs in the fall of 2011. Via this action one of the most important Noctuidae research material ever assembled by a private person in Hungary is going to be preserved in a public museum (Figs 2–3). The collection, which contains 141,000 specimens in 983 drawers, has two main parts. One is the Noctuidae collection of appr. 120,000 specimens in 800 drawers, accompanied by 12,000 genitalia slides and colour photographs of types and other historically important museum specimens with a similar number. This part of the collection not only contains a large number of primary types (Figs 4–5) but also secondary types ten times as many (Fig. 6). The other part is the Papilionoidea collection. This second part contains about 20,000 specimens in 183 drawers. Both parts include many exemplars in need of setting, labelling, sorting, and waiting for identification.

One of the conditions of his heirs for allowing the Hreblay collection to be deposited in a public museum was the publication of the collection's catalogue. This is a tremendous task and can be completed in steps planned for several years. Because of the large amount of type material the first step to be made was



Fig. 1. Márton Hreblay (circa in 1998) at his working-desk setting expedition material (photo: J. Heck)



Fig. 2. The authors are working in the room of the Hreblay collection pulling out drawers for databasing (January, 2013); the image shows approximately half of the entire collection (photo: Z. György)



Fig. 3. Drawers of the Hreblay collection displayed in the public exhibition “Leltárba véve – 9 643 086 tárgy, egy élő nemzeti természetrajzi gyűjtemény (9,643,086 specimens: catalogued – Hungary’s Living Natural History Collection)” between 18 May and 25 November, 2012 (photo: M. Katus)

the compilation of Márton Hreblay’s scientific works (BÁLINT *et al.* 2013). As a follow-up, the list of all the names proposed by Márton Hreblay for the newly discovered taxa could be collected, and the value of this contribution is enhanced by data of the primary types. This also determines which of his names are represented by primary types in his own collection.



Figs 4–5. *Harutaographa adusta* Hreblay et L. Ronkay, 1999: 4 = holotype in dorsal view; 5 = original labels (photos: G. Katona)



Fig. 6. Drawer no. 143 containing 265 paratype specimens of *Perigrapha scriptobella* Hreblay, 1996 (photo: Z. György)

The aim of this paper is to publish a list that contains all the names he described alone or with co-workers in scientific papers, book chapters or books. The list we present here is complete in this respect according to our best knowledge.

MATERIAL AND METHODS

Data on original descriptions, specimen depositories, type localities of primary types were compiled from the scientific papers, and recorded in MS Excel. The list and secondary information for tables were generated from this file.

Names are listed alphabetically by author(s) and year of publication. After the dash the original binomen is given with citation of the original paper and page number, where the name appears and has been described; after this the sex of the holotype specimen is indicated, followed by country of origin and (after the colon in quotation marks) the type locality taken verbatim from the original description; the depository of the holotype is the last information (in parentheses).

Primary type specimen depositories and their acronyms used in the present paper: BMNH = Natural History Museum, London, United Kingdom (formerly British Museum of Natural History); CCN = collection of Prof Dr Clas Naumann

deposited in MAK; CGR = private collection of Mr Gábor Ronkay (Budapest, Hungary); CGYF = private collection of Mr György Fábrián (Budapest, Hungary); CHH = private collection of Dr Hermann Hacker (Schwanfeld, Germany), affiliated with ZSM; CJP = collection of Mr Jacques Plante deposited in Muséum d'Histoire Naturelle, Geneva, Switzerland; CMF = private collection of Dr Michael Fibiger (Sorø, Denmark); CMH = collection of Dr Márton Hreblay deposited in HNHM; CPGY = private collection of Dr Péter Gyulai (Miskolc, Hungary); CSTK = private collection of Mr Sándor Tibor Kovács (Szeged, Hungary); CWS = private collection of Dr Wolfgang Speidel (München, Germany), affiliated with ZSM; HNHM = Hungarian Natural History Museum, Budapest, Hungary; HNHM-GR = specimen indicated to be deposited in HNHM via Gábor Ronkay; HNHM-PGY = specimen indicated to be deposited in HNHM via Péter Gyulai; IRSNB = Institut Royal des Sciences Naturelles de Belgique, Brussels, Belgium; MAK = Zoologisches Forschungsinstitut und Museum Alexander Koenig, Bonn, Germany; MNHU = Museum für Naturkunde der Humboldt-Universität, Berlin, Germany; NHMW = Naturhistorisches Museum, Wien, Austria; NIAES = Laboratory of Insect Systematics, National Institute of Agro-Environmental Sciences, Tsukuba, Japan; NMNS = National Museum of Natural Science, Taichung, Taiwan; NNM = Naturalis, National Museum of Natural History, Leiden, The Netherlands; NSMT = National Science Museum, Tokyo, Japan; UO = University of Osaka, College of Agriculture, Entomological Laboratory, Osaka, Japan; ZIN = Zoological Institute, Russian Academy of Sciences, Sankt-Petersburg, Russia; ZMH = Zoological Museum, University of Helsinki, Finland; ZMK = Zoologisk Museum, København, Denmark; ZSM = Zoologische Staatssammlung, München, Germany.

LIST OF TAXA DESCRIBED AS NEW BY MÁRTON HREBLAY

Species-group names

- abdita* Hreblay et Yoshimatsu, 1998 – Described as *Mythimna (Hyphilare) abdita* in HREBLAY *et al.* (1998a: 404). Holotype male, Indonesia: “Mt Goliath, Centr., Dutch N. Guinea” (BMNH).
- achrysa* L. Ronkay, Varga et Hreblay, 1998 – Described as *Polymixis achrysa* in RONKAY *et al.* (1998: 247). Holotype male, Turkmenistan: “USSR, Turkmenia, Kopet-Dagh Mts, Firyuza” (HNHM).
- actinea* Hreblay et L. Ronkay, 1999 – Described as *Euplexidia actinea* in HREBLAY & RONKAY (1999: 561). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 6 km W Cat Cat” (HNHM).
- aculeata* Hreblay et L. Ronkay, 1998 – Described as *Xanthia aculeata* in HREBLAY & RONKAY (1998: 243). Holotype male, Nepal: “Nepal, Ganesh Himal, 1 km E of Gadrang” (CMH).
- acuta* Hreblay et Plante, 1998 – Described as *Perissandria sheljuzhkoi acuta* in HREBLAY *et al.* (1998b: 125). Holotype male, China: “China, Tibet, Karo La (Pass), W side” (CJP).

- adusta* Hreblay et L. Ronkay, 1999 – Described as *Harutaographa adusta* in HREBLAY & RONKAY (1999: 492). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Inthanon” (CMH) (Figs 4–5).
- aestivalis* Hreblay et L. Ronkay, 1998 – Described as *Nyctycia aestivalis* in HREBLAY & RONKAY (1998: 202). Holotype female, Pakistan: “Pakistan, Himalaya Mts, Kaghan valley, 20 km NE Balakot, Tathabaya” (HMHM).
- akos* Hreblay, 1996 – Described as *Perigrapha (Harutaographa) akos* in HREBLAY (1996a: 81). Holotype male, Tajikistan: “USSR, Tadzhikistan, Hissarskij Mts, Romit” (ZMH).
- albicomma* L. Ronkay, Varga et Hreblay, 1998 – Described as *Haemosia albicomma* in RONKAY *et al.* (1998: 273). Holotype male, Turkmenistan: “Turkmenistan, Kopet-Dagh Mts, 5 km S of Chuli” (HNHM).
- albilabiata* Gyulai et Hreblay, 2001 – Described as *Trichanarta albilabiata* in GYULAI & HREBLAY (2001: 711). Holotype female, China: “Tibet West, Gandisa Mts, 100 km nord of Coqen” (HNHM-PGY).
- albimacula* Hreblay et L. Ronkay, 1998 – Described as *Oroplexia albimacula* in HREBLAY & RONKAY (1998: 256). Holotype male, China: “China, Tibet, 8 km S of Nyalam” (CMH).
- albiorbis* Hreblay et L. Ronkay, 1998 – Described as *Polymixis albiorbis* in HREBLAY & RONKAY (1998: 189). Holotype male, Nepal: “Nepal, Ganesh Himal, Kalchet, NE slope of Khurpudanda Pass” (CGR).
- albistigma* Hreblay et L. Ronkay, 1998 – Described as *Lithopolia albistigma* in HREBLAY & RONKAY (1998: 163). Holotype male, Taiwan: “Taiwan, Prov. Taitung, 2 km E of Hsiangyang” (CGYF).
- albistigma* Hreblay et L. Ronkay, 1998 – Described as *Loxopamea albistigma* in HREBLAY *et al.* (1998b: 146). Holotype male, China: [Ta-Tsien-Lou] (BMNH). Note: The original description states that the type locality is “Chasseurs Thibétains”, but actually it is Tat-Sien-Lou (present day Kangding, W. Sichuan, China) from where resident people supplied material for various European lepidopterists before the World War I.
- albitractora* Hreblay et Plante, 1995 – Described as *Loxopamea albitractora* in HREBLAY & PLANTE (1995b: 545). Holotype male, China: “Chine, Tibet, 8 km of Nyalam” (CMH).
- albivariiegata* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Nyctycia albivariiegata* in HREBLAY *et al.* (1999b: 20). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- albostrigata* Hreblay et Yoshimatsu, 1998 – Described as *Mythimna (Hyphilare) albostrigata* in HREBLAY *et al.* (1998a: 393). Holotype male, China: “[China], Kuatan” (MAK).
- albovenosa* Hreblay, 1999 – Described as *Mythimna (Hyphilare) albovenosa* in HREBLAY *et al.* (1999a: 382). Holotype male, India: “S-INDIA, Kerala, 7 km N Munnar, Eravikulam, NP” (CMH).
- albovitta* Hreblay et L. Ronkay, 2000 – Described as *Asidemia albovitta* in HREBLAY & RONKAY (2000: 31). Holotype male, Taiwan: “Taiwan, Prov. Hualien, Pilushernmuh” (NMNS).
- alfa* Hreblay et Plante, 1998 – Described as *Sympistis alfa* in HREBLAY *et al.* (1998b: 137). Holotype male, China: “China, Tibet, Karo La (Pass), W side” (CMH).
- alta* Hreblay et L. Ronkay, 1998 – Described as *Euxoa amorpha alta* in HREBLAY *et al.* (1998b: 113). Holotype male, China: “China, Tibet, 1 km S of Paiku Tso” (CMH).
- altiphila* Hreblay et Legrain, 1996 – Described as *Mythimna (Morphopoliana) altiphila* in HREBLAY (1996b: 147). Holotype male, Ethiopia: “Ethiopia, Addis Abeba” (HNHM).
- altissima* Hreblay et L. Ronkay, 1998 – Described as *Paraxestia altissima* in HREBLAY & RONKAY (1998: 133). Holotype male, Nepal: “Nepal, Annapurna Himal, Mesokantu Pass, SE slope” (CGR).

- angusta* Hreblay et L. Ronkay, 1998 – Described as *Haderonia praecipua angusta* in HREBLAY & RONKAY (1998: 150). Holotype male, Nepal: “Nepal, Annapurna Himal, 11 km SW of Jomsom, Noma pasture” (CGR).
- annamica* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Agrochola albirena annamica* in HREBLAY *et al.* (1999b: 43). Holotype male, Vietnam: “Vietnam N, Prov. Lao Cai, Sa Pa” (HNHM).
- annamica* Hreblay et L. Ronkay, 1999 – Described as *Owadaglea elongata annamica* in HREBLAY & RONKAY (1999: 547). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- annapurna* Hreblay et L. Ronkay, 1998 – Described as *Estimata annapurna* in HREBLAY & RONKAY (1998: 139). Holotype male, Nepal: “Nepal, Annapurna Himal, Mesokantu Pass” (CGR).
- annapurna* Hreblay et L. Ronkay, 1998 – Described as *Euplexia annapurna* in HREBLAY & RONKAY (1998: 262). Holotype female, Nepal: “Annapurna Himal, 10 km SE of Jomsom” (CGR).
- annapurnae* Hreblay et L. Ronkay, 1999 – Described as *Polymixis albiorbis annapurnae* in HREBLAY & RONKAY (1999: 548). Holotype male, Nepal: “Nepal, Annapurna Himal, 11 km SW of Jomsom, Noma pasture” (HNHM-GR).
- anonyma* Hreblay et Plante, 1996 – Described as *Hermonassa anonyma* in HREBLAY & PLANTE (1996: 663). Holotype male, Nepal: “Nepal, Ganesh Himal, 3 km SE from Somdang” (CMH).
- anonyma* Hreblay et L. Ronkay, 1998 – Described as *Conistra anonyma* in HREBLAY & RONKAY (1998: 230). Holotype male, Taiwan: “Taiwan, Prov. Kao-Hsiung, 15 km NE of Taoyuan” (CGYF).
- apameoides* Hreblay et L. Ronkay, 1998 – Described as *Oroplexia apameoides* in HREBLAY & RONKAY (1998: 257). Holotype male, Nepal: “Nepal, Lapchi Kang Range, 4 km SW of Tselaphu (Doupseyding)” (CMH).
- aquila* Hreblay et L. Ronkay, 1998 – Described as *Xestia aquila* in HREBLAY & RONKAY (1998: 137). Holotype male, Nepal: “Nepal, Ganesh Himal, 3 km SE from Somdang” (CMH).
- arcanus* Hreblay et Plante, 1995 – Described as *Sideridis (Dianthivora) arcanus* in HREBLAY & PLANTE (1995a: 137). Holotype male, Nepal: “Népal, Ganesh Himal, 3 km NE of Sunpati” (CMH).
- archar* L. Ronkay, Varga et Hreblay, 1998 – Described as *Agrochola oropotamica archar* in RONKAY *et al.* (1998: 256). Holotype male, Turkmenistan: “USSR, Turkmenia, 80 km SE of Tedjen” (CGR).
- ardescentina* Hreblay et L. Ronkay, 1998 – Described as *Conistra ardescentina* in HREBLAY & RONKAY (1998: 231). Holotype female, Nepal: “Ganesh Himal, near Godlang” (CGR).
- argentata* Hreblay et Yoshimatsu, 1998 – Described as *Mythimna (Hyphilare) argentata* in HREBLAY *et al.* (1998a: 406). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 3 km E of Tili” (CMH).
- argentifera* Hreblay, 1998 – Described as *Mythimna (Hyphilare) argentifera* in HREBLAY *et al.* (1998a: 407). Holotype male, Indonesia: “Fort de Nock, [Sumatra]” (NNM).
- argillosa* Hreblay et L. Ronkay, 1998 – Described as *Agrotis argillosa* in HREBLAY *et al.* (1998b: 120). Holotype male, China: “Gyantse” (MNHU).
- aristata* Hreblay et L. Ronkay, 1995 – Described as *Himalistra aristata* in HREBLAY *et al.* (1995: 61). Holotype male, Nepal: “Nepal, Langtang, between Cholang Pati and Dimsa” (HNHM).
- ariuna* Hreblay, 1991 – Described as *Orthosia (Orthosia) ariuna* in HREBLAY (1991: 195). Holotype male, Mongolia: “Mongolia, Bulgan aimak, 64 km W of Erdenecant” (CMH).
- armata* Hreblay et L. Ronkay, 1998 – Described as *Meganyctycia armata* in HREBLAY & RONKAY (1998: 207). Holotype male, Nepal: “Nepal, Arun valley, 12 km N of Chitre” (CMH).
- arnyekolta* L. Ronkay, Varga et Hreblay, 1998 – Described as *Autophila luxuriosa arnyekolta* in RONKAY *et al.* (1998: 277). Holotype male, Turkmenistan: “Turkmenistan, Kopet-Dagh Mts, 6 km S of Ipay-Kala” (HNHM).

- atrifusa* Hreblay et Kobayashi, 1997 – Described as *Telorta atrifusa* in HREBLAY & RONKAY (1997: 56). Holotype female, Taiwan: “Taiwan, Prov. Nantou, Tayuling” (CGYF).
- asymmetrica* Hreblay et L. Ronkay, 1998 – Described as *Nyctycia asymmetrica* in HREBLAY & RONKAY (1998: 201). Holotype male, Nepal: “Nepal, Arun valley: 12 km N of Hille” (CMH).
- atra* Hreblay et L. Ronkay, 1998 – Described as *Odontestera atra* in HREBLAY & RONKAY (1998: 149). Holotype male, Nepal: “Nepal, Annapurna Himal, valley of Kali Gandaki, near Ghasa” (CGR).
- atra* Hreblay et L. Ronkay, 1998 – Described as *Xenotrachea atra* in HREBLAY & RONKAY (1998: 268). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, 10 km S of Lukla, Bupsa” (CMH).
- atra* Hreblay et Gyulai, 1998 – Described as *Isochlora atra* in HREBLAY *et al.* (1998b: 110). Holotype male, China: “China, Xin-Jiang, Altun shan” (CPGY).
- atrata* Hreblay et L. Ronkay, 1999 – Described as *Orthosia (Euchoristea) limbata atrata* in HREBLAY & RONKAY (1999: 499). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Inthanon” (CMH).
- attika* L. Ronkay, Yela & Hreblay, 2001 – Described as *Leucochlaena (Leucochlaena) muscosa attika* in RONKAY *et al.* (2001: 256). Holotype male, Greece: “Attiki, Erythrae” (CMF).
- attila* Hreblay et L. Ronkay, 1999 – Described as *Agrochola attila* in HREBLAY & RONKAY (1999: 527). Holotype male, Thailand: “Thailand, Changwat Nan, 30 km E of Pua” (CMH).
- attila* Hreblay et L. Ronkay, 1999 – Described as *Nyctycia attila* in HREBLAY & RONKAY (1999: 537). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, 20 km NW of Sop Kha, 2 km S of Kop Dong” (CMH).
- attila* Hreblay et L. Ronkay, 2000 – Described as *Odontestera attila* in HREBLAY & RONKAY (2000: 13). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, 2 km S of Ban Kum” (CMH).
- aulombardi* Hreblay, 1992 – Described as *Conistra aulombardi* in HREBLAY (1992b: 542). Holotype male, Pakistan: “Pakistan, Baltistan, Lac. Satpara, 8 km sud de Skardu” (CJP).
- aurantiaca* Hreblay et Plante, 1995 – Described as *Valeriodes aurentiaca* in HREBLAY & PLANTE (1995a: 138). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, Lukla” (CJP).
- aurotrigoni* Hreblay, 1998 – Described as *Mythimna (Mythimna) aurotrigoni* in HREBLAY *et al.* (1998a: 385). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok” (CMH).
- auroviridis* Hreblay et L. Ronkay, 1999 – Described as *Valeriodes auroviridis* in HREBLAY & RONKAY (1999: 558). Holotype male, Nepal: “East-Nepal, Deorali Danda, Tseram” (CMH).
- averna* Hreblay et L. Ronkay, 1997 – Described as *Amphipyra averna* in HREBLAY & RONKAY (1997: 81). Holotype female, Taiwan: “Taiwan, Prov. Taoyuan, 16 km E of Fuhsing” (CMH).
- barna* Hreblay et L. Ronkay, 1998 – Described as *Owadaglaea barna* in HREBLAY & RONKAY (1998: 227). Holotype female, Nepal: “Ganesh Himal, 1 km E of Gadrang” (CMH).
- basivittata* L. Ronkay, Varga et Hreblay, 1998 – Described as *Cryphia basivittata* in RONKAY *et al.* (1998: 261). Holotype male, Turkmenistan: “Turkmenistan, Kopet-Dagh Mts, Dushak Mt” (HNHM).
- bea* Hreblay et Plante, 1998 – Described as *Euchalcia bea* in HREBLAY *et al.* (1998b: 154). Holotype male, China: “China, Tibet, 50 km N of Nyalam” (CMH).
- beata* Hreblay et L. Ronkay, 1998 – Described as *Polymixis beata* in HREBLAY & RONKAY (1998: 188). Holotype female, Nepal: “Nepal, Annapurna Himal, 5 km SE of Jomsom, Thadung valley” (CGR).
- behouneki* Hreblay et Plante, 1995 – Described as *Lacanobia behouneki* in HREBLAY & PLANTE (1995a: 137). Holotype male, Nepal: “Nepal, Ganesh Himal, 3 km NE of Sunpati” (CMH).

- belastigma* Hreblay et L. Ronkay, 1998 – Described as *Trachea belastigma* in HREBLAY & RONKAY (1998: 266). Holotype male, Nepal: “Nepal, Annapurna Himal, Banthanti” (CGR).
- benescripta* Hreblay et L. Ronkay, 1995 – Described as *Estagrotis benescripta* in HREBLAY & RONKAY (1995a: 244). Holotype male, Nepal: “Nepal, Annapurna Himal, near Deorali” (HNHM).
- beri* Gyulai et Hreblay, 1992 – Described as *Oncocnemis beri* in GYULAI *et al.* (1992: 39). Holotype male, Uzbekistan: “USSR, Uzbekistan SSR, W Tien Shan, Mts Chimgan” (CMH).
- beta* Hreblay et Plante, 1998 – Described as *Sympistis beta* in HREBLAY *et al.* (1998b: 138). Holotype male, China: “China, Tibet, 20 km S of Saga, 2 km S of Lodu Loney Tso” (CMH).
- bicolor* Hreblay et L. Ronkay, 1995 – Described as *Dasypolia (Dasypolia) bicolor* in HREBLAY & RONKAY (1995b: 368). Holotype male, China: “China, Tibet: 4 km S of Karru-Ochen Tso” (HNHM).
- bicolorata* Hreblay et L. Ronkay, 1998 – Described as *Harutaographa bicolorata* in HREBLAY & RONKAY (1998: 155). Holotype male, Nepal: “Nepal, Ganesh Himal, 1 km E of Gadrang” (CMH).
- bidui* Hreblay et Plante, 1996 – Described as *Perigrapha (Harutaographa) bidui* in HREBLAY (1996a: 79). Holotype female, Pakistan: “N-Pakistan, 5 km S of Rattu” (CMH).
- biformis* Hreblay et Kononenko, 1999 – Described as *Diphtherocome biformis* in HREBLAY & KONONENKO (1999: 678). Holotype male, China: “[China, Prov. Shensi], Tapaishan im Tsi-ling, Sued-Shenshi” (MAK).
- bifurcata* Hreblay et Plante, 1996 – Described as *Oroplexia bifurcata* in HREBLAY & PLANTE (1996: 666). Holotype male, India: “N. India, Himalaya Mts, Maj” (ZSM).
- bifurcata* Hreblay et L. Ronkay, 1998 – Described as *Xestia bifurcata* in HREBLAY & RONKAY (1998: 135). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, 7 km E of Lukla” (CMH).
- bifurcata* Hreblay et Plante, 1998 – Described as *Athetis bifurcata* in HREBLAY *et al.* (1998b: 151). Holotype male, China: “China, Tibet, 10 km E of Nyemo” (CMH).
- bigu* Hreblay et L. Ronkay, 1999 – Described as *Diphtherocome bigu* in HREBLAY & RONKAY (1999: 566). Holotype male, Nepal: “Nepal, Mt Kalinchok, 5 km W of Bigu” (CMH).
- biscornuta* Hreblay, Legrain et Yoshimatsu, 1998 – Described as *Mythimna (Hyphilare) biscornuta* in HREBLAY *et al.* (1998a: 394). Holotype male, Indonesia: “[Indonesia] Sumatra, Prapat HW” (BMNH).
- bodii* Hreblay et L. Ronkay, 1998 – Described as *Lithopolia bodii* in HREBLAY & RONKAY (1998: 162). Holotype male, Nepal: “Nepal, Dhaulagiri Himal, 2 km S of Lebang” (CMH).
- brahma* Hreblay et L. Ronkay, 1998 – Described as *Harutaographa brahma* in HREBLAY & RONKAY (1998: 156). Holotype male, Nepal: “Nepal, Ganesh Himal, 2 km W of Thangjet” (CMH).
- brevica* Hreblay et Legrain, 1998 – Described as *Mythimna (Hyphilare) brevica* in HREBLAY *et al.* (1998a: 402). Holotype male, Papua New Guinea: “New Guinea, Highlands, Mt Michael, Frigano” (BMNH).
- brevicornata* Hreblay et L. Ronkay, 1999 – Described as *Meganycytia brevicornata* in HREBLAY & RONKAY (1999: 542). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 7 km W Sa Pa” (HNHM).
- brevipennata* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Daseuplexia brevipennata* in HREBLAY *et al.* (1999b: 33). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 3 km NW Cat Cat” (HNHM).
- bryochlora* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Diphtherocome bryochlora* in HREBLAY *et al.* (1999b: 83). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 3 km NW Cat Cat” (HNHM).

- bugeri* Hreblay et Thöny, 1995 – Described as *Checupa bugeri* in HREBLAY & THÖNY (1995: 27). Holotype male, Philippines: “Philippinen, Mindanao, S-Mt Kitanglad” (CMH).
- cadoui* Hreblay et L. Ronkay, 1998 – Described as *Neurois cadoui* in HREBLAY & RONKAY (1998: 144). Holotype male, Nepal: “Nepal, E., Chagam” (HNHM).
- caeca* Hreblay et L. Ronkay, 1997 – Described as *Polia mortua caeca* in HREBLAY & RONKAY (1997: 28). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 7 km SW of Tayüling” (CMH).
- caelestissima* Hreblay et L. Ronkay, 1998 – Described as *Euchalcia caelestissima* in HREBLAY *et al.* (1998b: 153). Holotype male, China: “China, Tibet, 50 km N of Nyalam” (CMH).
- caesia* Hreblay et Plante, 1994 – Described as *Himalistra caesia* in HREBLAY *et al.* (1994: 259). Holotype female, Pakistan: “N-Pakistan, 10 km SW of Astor, Rama” (CJP).
- caesia* Hreblay, 1998 – Described as *Apamea caesia* in HREBLAY & RONKAY (1998: 248). Holotype male, Nepal: “Ganesh Himal, 2 km W of Thangjet” (CMH).
- caliginata* Hreblay, 1999 – Described as *Mythimna (Hyphilare) ferrago caliginata* in HREBLAY *et al.* (1999a: 379). Holotype male, Pakistan: “Pakistan, Prov. Jammu & Kaschmir, Gilgit valley, Jundrot village” (CMH).
- callilhasa* Hreblay et L. Ronkay, 1998 – Described as *Agrotis callilhasa* in HREBLAY *et al.* (1998b: 118). Holotype male, China: “Tibet South, Nyainqentanglha Mts, Lhasa Nord” (CPGY).
- cameroni* Hreblay et Yoshimatsu, 1998 – Described as *Mythimna (Mythimna) cameroni* in HREBLAY *et al.* (1998a: 386). Holotype male, Malaysia: “Malaysia, Pahang state, Cameron Highland am Tanah Rata” (CGR).
- caovansungi* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Isolasia caovansungi* in HREBLAY *et al.* (1999b: 28). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- cara* Hreblay et L. Ronkay, 1998 – Described as *Xestia cara* in HREBLAY & RONKAY (1998: 136). Holotype male, Nepal: “Nepal, Annapurna Himal, 10 km SE of Jomsom” (CGR).
- cbgurungi* Hreblay et L. Ronkay, 1998 – Described as “*Mniotype*” *cbgurungi* in HREBLAY & RONKAY (1998: 192). Holotype male, Nepal: “Nepal, Tanahoun distr., Bimlanager village” (HNHM).
- celebina* Hreblay, 1998 – Described as *Mythimna (Hyphilare) celebina* in HREBLAY *et al.* (1998a: 403). Holotype male, Indonesia: “S-Sulawesi, (Celebes), Puncak Palopo” (CMH).
- cerriformis* Hreblay et L. Ronkay, 1998 – Described as *Dryobotodes cerriformis* in HREBLAY & RONKAY (1998: 184). Holotype male, Nepal: “Nepal, Langtang, near Chandrabari” (HNHM).
- chapa* Hreblay, 1999 – Described as *Analetia (Anapoma) chapa* in HREBLAY *et al.* (1999a: 388). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok, 16 km NW of Fang” (CMH).
- chengai* Hreblay et L. Ronkay, 1998 – Described as *Trichoridia chengai* in HREBLAY & RONKAY (1998: 177). Holotype male, Nepal: “Nepal, Lapchi Kang Range, 4 km SW of Tselaphu (Doupseyding)” (CMH).
- chhiringi* Hreblay, 1998 – Described as *Apamea chhiringi* in HREBLAY & RONKAY (1998: 249). Holotype male, Nepal: “Ganesh Himal, 3 km NE of Sunpati” (CMH).
- chiangmai* Hreblay et Yoshimatsu, 1998 – Described as *Mythimna (Mythimna) chiangmai* in HREBLAY *et al.* (1998a: 387). Holotype male, Thailand: “Thailand, Changwat Mae Hong Song, 1 km S of Bahundanda” (CMH).
- chlorata* Hreblay et L. Ronkay, 1999 – Described as *Isolasia chlorata* in HREBLAY & RONKAY (1999: 541). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok, 18 km NW of Fang” (CMH).
- chlorobesa* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Mniotype chlorobesa* in HREBLAY *et al.* (1999b: 65). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).

- chloromagna* Hreblay et L. Ronkay, 1998 – Described as *Daseuplexia chloromagna* in HREBLAY & RONKAY (1998: 211). Holotype female, Nepal: “Annapurna Himal, 1 km E of Gorepani” (CMH).
- chloronympha* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Diphtherocome chloronympha* in HREBLAY et al. (1999b: 85). Holotype male, Nepal: “Nepal, Koshi, Terhathum area, Tinjure Phedi” (CGR).
- cineracea* Hreblay et L. Ronkay, 1998 – Described as *Potnyctycia cineracea* in HREBLAY & RONKAY (1998: 204). Holotype female, Nepal: “Ganesh Himal, 2 km W of Gholjong” (CGR).
- cinerea* Hreblay et L. Ronkay, 1998 – Described as *Harutaeographa cinerea* in HREBLAY & RONKAY (1998: 155). Holotype female, Nepal: “Nepal, Ganesh Himal, near Slya” (CGR).
- citrina* Hreblay et L. Ronkay, 1998 – Described as *Apostema citrina* in HREBLAY & RONKAY (1998: 194). Holotype female, Nepal: “Nepal, Langtang, near Chandrabari” (CGR).
- claricostata* Hreblay et Plante, 1995 – Described as *Hermonassa claricostata* in HREBLAY & PLANTE (1995a: 135). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, Lamjura Pass” (CJP).
- clarior* Hreblay, 1992 – Described as *Mythimna (Aletia) clarior* in HREBLAY (1992a: 520). Holotype male, Mongolia: “Mongolei, Chovd aimak, Mongol. Altaj Gebirge, Uljasutajn gol, 45 km NNO von Somon Bulgan” (HNHM).
- clavata* Hreblay et L. Ronkay, 1999 – Described as *Mniotype clavata* in HREBLAY & RONKAY (1999: 550). Holotype male, Nepal: “East-Nepal, Surke Danda, 2 km NE of Suketar” (CMH).
- complicata* Hreblay, 1999 – Described as *Analetia (Anapoma) complicata* in HREBLAY et al. (1999a: 389). Holotype male, Vietnam: “N-Vietnam, Mt Fan-si-pan, Cha-pa” (CMH).
- confusa* Hreblay et L. Ronkay, 1995 – Described as *Dasytopia (s. l.) confusa* in HREBLAY & RONKAY (1995b: 370). Holotype male, China: “China, Tibet: 4 km S of Karru-Ochen Tso” (HNHM).
- conistroides* Hreblay et L. Ronkay, 1995 – Described as *Dasytopia (s. l.) conistroides* in HREBLAY & RONKAY (1995b: 373). Holotype female, Nepal: “Nepal. Ganesh Himal, 3 km SE from Somdang” (HNHM).
- conjunctura* Hreblay et L. Ronkay, 1998 – Described as *Oroplexia conjunctura* in HREBLAY & RONKAY (1998: 255). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, 7 km E of Lukla” (CMH).
- consimilis* Hreblay et L. Ronkay, 1998 – Described as *Nyctycia consimilis* in HREBLAY & RONKAY (1998: 200). Holotype male, Nepal: “Ganesh Himal, 2 km W of Thangjet” (CMH).
- conversa* Hreblay et Yoshimatsu, 1998 – Described as *Mythimna (Hyphilare) conversa* in HREBLAY et al. (1998a: 408). Holotype male, Indonesia: “Sumatra, Prapat” (BMNH).
- copria* Hreblay et Plante, 1995 – Described as *Diarsia copria* in HREBLAY & PLANTE (1995b: 539). Holotype male, Nepal: “Nepal, Ganesh Himal, 1 km E of Yurekharka” (CMH).
- cornutifera* Hreblay et L. Ronkay, 1997 – Described as *Chortodes cornutifera* in HREBLAY & RONKAY (1997: 61). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 7 km SW of Tayiling” (CMH).
- coronata* Hreblay et L. Ronkay, 1999 – Described as *Xestia coronata* in HREBLAY & RONKAY (1999: 487). Holotype male, Nepal: “East-Nepal, Deorali Danda, Tseram” (CMH).
- costigera* Hreblay et L. Ronkay, 1998 – Described as *Hemiglaea costigera* in HREBLAY & RONKAY (1998: 220). Holotype male, Nepal: “Nepal, Arun valley, 21 km N of Hille” (CMH).
- criaquatica* Hreblay et L. Ronkay, 1999 – Described as *Lithopolia (Hiemopolia) criaquatica* in HREBLAY & RONKAY (1999: 513). Holotype male, Nepal: “Nepal, Koshi, Terhathum area, above Gorja, Tsishopani” (HNHM).
- crisifera* Hreblay et L. Ronkay, 1997 – Described as *Potnyctycia crisifera* in HREBLAY & RONKAY (1997: 52). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 5 km SW of Tayiling” (CGYF).
- cuprea* Hreblay et L. Ronkay, 1998 – Described as *Eupsilia cuprea* in HREBLAY & RONKAY (1998: 183). Holotype female, Nepal: “Ganesh Himal, between Godlang and Nesim” (CGR).

- curta* Hreblay, 1998 – Described as *Mythimna (Hyphilare) simplex curta* in HREBLAY *et al.* (1998a: 399). Holotype male, Philippines: “Philippinen, Mindanao, S-Mt Kitanglad” (CMH).
- cyanochlora* Hreblay et L. Ronkay, 1998 – Described as *Mniotype cyanochlora* in HREBLAY & RONKAY (1998: 191). Holotype male, Nepal: “Nepal, 4 km SW of Kalinchok peak” (CMH).
- csaba* Hreblay et L. Ronkay, 1999 – Described as *Lithopolia (Hiemopolia) csaba* in HREBLAY & RONKAY (1999: 511). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok, 16 km NW of Fang” (CMH).
- csanadii* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Mniotype csanadii* in HREBLAY *et al.* (1999b: 73). Holotype male, Nepal: “Nepal, Langtang, near Chandrabari” (HNHM).
- csoevarii* Hreblay et L. Ronkay, 2000 – Described as *Xestia fuscostigma csoevarii* in HREBLAY & RONKAY (2000: 4). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 3 km SW of Tsuifeng” (CMH).
- csorbagabori* L. Ronkay, Varga et Hreblay, 1998 – Described as *Polymixis csorbagabori* in RONKAY *et al.* (1998: 252). Holotype male, Turkey: “Turkey, Prov. Van, Kuzgunkiran Gecidi” (HNHM).
- csorbai* Hreblay et L. Ronkay, 1998 – Described as *Mniotype csorbai* in HREBLAY & RONKAY (1998: 193). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Sa Pa” (HNHM).
- daemon* Hreblay et Legrain, 1996 – Described as *Mythimna (Mythimna) daemon* in HREBLAY *et al.* (1996: 95). Holotype male, Nepal: “Nepal, Annapurna Himal, Thini, 1 km S of Jomson” (CGR).
- dagestana* Hreblay et L. Ronkay, 1998 – Described as *Isochlora dagestana* in HREBLAY *et al.* (1998b: 107). Holotype male, Russia: “(S-Russia), Dagestan, Fl. Samur” (CMH).
- dailekhi* Hreblay et L. Ronkay, 1999 – Described as *Lithophane dailekhi* in HREBLAY & RONKAY (1999: 523). Holotype female, Nepal: “West-Nepal, 11 km N of Dailekh” (CMH).
- daliangi* Hreblay et L. Ronkay, 1998 – Described as *Diptherocome daliangi* in HREBLAY *et al.* (1998b: 111). Holotype male, China: “China, Setchuan, Daliang Shan” (CMH).
- decoronata* Hreblay, 1996 – Described as *Analetia (Anapoma) decoronata* in HREBLAY *et al.* (1996: 117). Holotype male, Nepal: “Nepal, Langtang 5 km NNE of Dhunche, Barkhu” (CMH).
- deletaiwana* Hreblay et L. Ronkay, 1998 – Described as *Amphipyra deletaiwana* in HREBLAY & RONKAY (1998: 275). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 1 km W of Tatachia peak” (CMH).
- delineata* Hreblay et L. Ronkay, 1998 – Described as *Dasytopia delineata* in HREBLAY & RONKAY (1998: 187). Holotype female, Nepal: “Nepal, Kalinchok area, 6 km SW of Kalinchok peak” (CMH).
- denticula* Hreblay et L. Ronkay, 1999 – Described as *Trichoridia denticula* in HREBLAY & RONKAY (1999: 515). Holotype male, Nepal: “Nepal, Ganesh Himal, Jageswar Kund” (CMH).
- diffusa* Hreblay et L. Ronkay, 1998 – Described as *Conistra aulombardi diffusa* in HREBLAY & RONKAY (1998: 231). Holotype male, Nepal: “Ganesh Himal, 2 km W of Thangjet” (CMH).
- dilikoti* Hreblay et L. Ronkay, 1999 – Described as *Estagrotis dilikoti* in HREBLAY & RONKAY (1999: 533). Holotype female, Nepal: “West-Nepal, 21 km N of Dailekh” (CMH).
- diluta* L. Ronkay, Varga et Hreblay, 1998 – Described as *Bornolis crinomima diluta* in RONKAY *et al.* (1998: 254). Holotype male, Turkmenistan: “USSR, Turkmenia, Kopet-Dagh Mts, 25 km E of Nochkur, Karayalchi valley” (HNHM).
- dilutissima* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Isolasia dilutissima* in HREBLAY *et al.* (1999b: 32). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- dimorpha* Hreblay et L. Ronkay, 1995 – Described as *Himalistra dimorpha* in HREBLAY & RONKAY (1995a: 238). Holotype male, Nepal: “Nepal, Ganesh Himal, 2 km E of Yurekharka” (HNHM).

- discestroides* Hreblay et L. Ronkay, 1998 – Described as *Pygmeopolia discestroides* in HREBLAY & RONKAY (1998: 161). Holotype male, Nepal: “Nepal, Ganesh Himal, 1 km E of Gadrang” (CMH).
- discoidea* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Rhynchaglaea discoidea* in HREBLAY et al. (1999b: 50). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- disseminata* Hreblay et Plante, 1995 – Described as *Xenotrachea disseminata* in HREBLAY & PLANTE (1995b: 544). Holotype male, Nepal: “Nepal, Taplejung area, Lal Kharka” (CMH).
- dissimilis* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Nyctycia dissimilis* in HREBLAY et al. (1999b: 11). Holotype male, Thailand: “Thailand, Doi Inthanon” (CHH).
- distincta* Hreblay et L. Ronkay, 1999 – Described as *Lithopolia (Pectinipolia) distincta* in HREBLAY & RONKAY (1999: 514). Holotype male, Thailand: “Thailand, Doi Inthanon” (CHH).
- diverticulata* Hreblay et Kononenko, 1999 – Described as *Diphtherocome diverticulata* in HREBLAY & KONONENKO (1999: 673). Holotype male, China: “[China, Prov. Yunnan], Likiang, Prov. Nord-Yuennan” (MAK).
- divida* Hreblay et L. Ronkay, 1998 – Described as *Hermonassa divida* in HREBLAY & RONKAY (1998: 132). Holotype male, Nepal: “Nepal, Ganesh Himal, 1 km E of Yurekharka” (CMH).
- doiinthoni* Hreblay et L. Ronkay, 1999 – Described as *Agrochola albirena doiinthoni* in HREBLAY & RONKAY (1999: 526). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Inthanon” (CMH).
- dora* Hreblay, 1992 – Described as *Conistra dora* in HREBLAY (1992b: 543). Holotype male, Turkmenistan: “Turkmenia, Kopet-Dagh Mts, 15 km s of Nochur” (CMH).
- dorsivitta* Hreblay et Plante, 1995 – Described as *Paraxestia dorsivitta* in HREBLAY & PLANTE (1995b: 542). Holotype male, Nepal: “Nepal, Ganesh Himak, Khurpudanda Pass” (CJP).
- duplicata* Hreblay et L. Ronkay, 1998 – Described as *Daseuplexia duplicata* in HREBLAY & RONKAY (1998: 211). Holotype male, Nepal: “Nepal, Ganesh Himal, between Godlang and Nesim” (CGR).
- dushaki* L. Ronkay, Varga et Hreblay, 1998 – Described as *Dasypteria templi dushaki* in RONKAY et al. (1998: 233). Holotype male, Turkmenistan: “USSR, Turkmenia, Kopet-Dagh Mts, 15 km SE of Nochur” (CMH).
- duskeimima* L. Ronkay, Varga et Hreblay, 1998 – Described as *Cryphia duskeimima* in RONKAY et al. (1998: 263). Holotype male, Turkmenistan: “Turkmenistan, Kopet-Dagh Mts, 6 km S of Ipay-Kala” (CMH).
- ebria* Hreblay et L. Ronkay, 1999 – Described as *Kisegira ebria* in HREBLAY & RONKAY (1999: 507). Holotype male, Thailand: “Thailand, Changwat Nan, 30 km E of Pua” (CMH).
- echinata* Hreblay et L. Ronkay, 1998 – Described as *Dasypteria (Dasymixis) echinata* in HREBLAY & RONKAY (1998: 187). Holotype male, Nepal: “Nepal, Ganesh Himal, Khurpudanda Pass” (CMH).
- elegantula* Hreblay et Plante, 1995 – Described as *Chandata elegantula* in HREBLAY & PLANTE (1995b: 543). Holotype male, Nepal: “Nepal, Taplejung area, Tambowa” (CJP).
- elongata* Hreblay et L. Ronkay, 1998 – Described as *Owadaglaea elongata* in HREBLAY & RONKAY (1998: 229). Holotype female, Nepal: “Nepal, Annapurna Himal, vic. Lume” (CGR).
- elphinia* Hreblay et L. Ronkay, 1999 – Described as *Harutaeographa elphinia* in HREBLAY & RONKAY (1999: 490). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 7 km W Sa Pa” (HNHM-GR).
- erecta* Hreblay, 1999 – Described as *Leucania erecta* in HREBLAY et al. (1999a: 385). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, 10 km NW of Fang” (CMH).

- erythrographa* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Apamea erythrographa* in HREBLAY *et al.* (1999b: 76). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 3 km NW Cat Cat” (HNHM).
- eugrapha* Hreblay et L. Ronkay, 1998 – Described as *Xestia eugrapha* in HREBLAY & RONKAY (1998: 137). Holotype male, Nepal: “Nepal, Annapurna Himal, 11 km SW of Jomsom, Noma pasture” (CGR).
- eupepla* L. Ronkay, Varga et Hreblay, 1998 – Described as *Conisania vidua eupepla* in RONKAY *et al.* (1998: 221). Holotype male, Turkmenistan: “Turkmenistan, Kopet-Dagh Mts, 6 km S of Ipay-Kala” (HNHM).
- euphemia* Hreblay et L. Ronkay, 1999 – Described as *Kisegira euphemia* in HREBLAY & RONKAY (1999: 506). Holotype male, Nepal: “Nepal, Koshi, Terhathum area, above Gorja, Tsisho-pani” (HNHM-GR).
- euplexina* Hreblay et L. Ronkay, 1998 – Described as *Blepharomima euplexina* in HREBLAY & RONKAY (1998: 213). Holotype male, Nepal: “Nepal, Arun valley, 12 km N of Hille” (CMH).
- euxoides* Hreblay et Plante, 1995 – Described as *Hermonassa euxoides* in HREBLAY & PLANTE (1995a: 136). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, Tragsindha Pass” (CJP).
- euxoides* Hreblay et L. Ronkay, 1998 – Described as *Conisania euxoides* in HREBLAY *et al.* (1998b: 129). Holotype male, China: “China, Tibet, 20 km S of Saga, 2 km S of Lodu Loney Tso” (CMH).
- excelsa* Hreblay et L. Ronkay, 1998 – Described as *Diarsia excelsa* in HREBLAY & RONKAY (1998: 143). Holotype male, Nepal: “Nepal, Kalinchok area, 4 km SW of Kalinchok peak” (CMH).
- exornata* Hreblay et L. Ronkay, 1997 – Described as *Panolis pinicortex exornata* in HREBLAY & RONKAY (1997: 28). Holotype male, Taiwan: “Taiwan, Prov Miaoli, 49 km E of Tungshih” (CMH).
- expallida* Hreblay et L. Ronkay, 1998 – Described as *Owadaglaea expallida* in HREBLAY & RONKAY (1998: 226). Holotype male, Nepal: “Nepal, Arun valley, 12 km N of Hille” (CMH).
- expatria* Hreblay et Plante, 1995 – Described as *Hermonassa expatria* in HREBLAY & PLANTE (1995b: 541). Holotype male, Nepal: “Nepal, Ganesh Himal, 2 km W of Thangjet” (CMH).
- expulsa* Hreblay, 1996 – Described as *Mythimna (Morphopolianna) expulsa* in HREBLAY (1996b: 138). Holotype male, Australia: “Australia, Mackay, Queensland” (BMNH).
- fabiani* Hreblay et L. Ronkay, 1998 – Described as *Perigrapha (Anorthoa) fabiani* in HREBLAY & RONKAY (1998: 152). Holotype male, Taiwan: “Taiwan, Prov. Taichung, 5 km NE of Tungshih, Wushihken” (CGYF).
- fabiani* Hreblay et L. Ronkay, 1998 – Described as *Oroplexia fabiani* in HREBLAY & RONKAY (1998: 253). Holotype male, Nepal: “Ganesh Himal, Nesukharka, 12 km S of Somdang” (CGYF).
- fabiani* L. Ronkay, Varga et Hreblay, 1998 – Described as *Polymixis fabiani* in RONKAY *et al.* (1998: 244). Holotype male, Pakistan: “Pakistan, Hindukush Mts, Shandur Pass, E slope” (CGYF).
- fakosharga* Hreblay et L. Ronkay, 1998 – Described as *Xestia fakosharga* in HREBLAY & RONKAY (1998: 138). Holotype male, Nepal: “Nepal, Langtang, 3 km SE of Syabru” (CGR).
- falcifera* Hreblay et L. Ronkay, 1999 – Described as *Mniotype falcifera* in HREBLAY & RONKAY (1999: 549). Holotype male, Nepal: “East-Nepal, Surke Danda, 2 km NE of Suketar” (CMH).
- fansipana* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Meganyctycia fansipana* in HREBLAY *et al.* (1999b: 25). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 3 km NW Cat Cat” (HNHM).
- faqiri* Hreblay et Plante, 1994 – Described as *Orthosia (Orthosia) faqiri* in HREBLAY & PLANTE (1994: 24). Holotype male, Pakistan: “NW-Pakistan, Kalam” (CMH).
- fatima* Hreblay, 1994 – Described as *Fatima fatima* in HREBLAY (1994: 245). Holotype male, Turkey: “Turkey, Prov. Maras, 1 km. N of Kahramanmaras” (CMH).

- fedra* Hreblay et Plante, 1994 – Described as *Orthosia (Orthosia) fedra* in HREBLAY & PLANTE (1994: 25). Holotype male, Pakistan: “N-Pakistan, 10 km SW of Astor, Rama” (CMH).
- felicita* Hreblay et Plante, 1995 – Described as *Hermonassa felicita* in HREBLAY & PLANTE (1995b: 541). Holotype male, Nepal: “Nepal, Taplejung area, 1 km of Suketar” (CJP).
- ferroboscuro* Hreblay et L. Ronkay, 1998 – Described as *Lasianobia ferroboscuro* in HREBLAY et al. (1998b: 131). Holotype male, China: “Chigatse” (MNHU).
- ferruginea* Hreblay et L. Ronkay, 1998 – Described as *Oroplexia ferruginea* in HREBLAY & RONKAY (1998: 258). Holotype male, Nepal: “Nepal, Annapurna Himal, Nangethanti” (HNHM).
- fibigeri* Hreblay et L. Ronkay, 1998 – Described as *Lithophane venusta fibigeri* in HREBLAY & RONKAY (1998: 181). Holotype male, Nepal: “Ganesh Himal, near Godlang” (HNHM).
- ficta* Hreblay et Plante, 1998 – Described as *Perissandria ficta* in HREBLAY et al. (1998b: 125). Holotype male, China: “China, Tibet, 10 km E of Tsou La (Pass)” (CJP).
- firyuza* L. Ronkay, Varga et Hreblay, 1998 – Described as *Eicomorpha firyuza* in RONKAY et al. (1998: 218). Holotype male, Turkmenistan: “USSR, Turkmenistan, Kopet-Dagh Mts, Firyuza” (CMH).
- fiyu* Hreblay et Yoshimatsu, 1998 – Described as *Leucania fiyu* in HREBLAY et al. (1998a: 413). Holotype male, Fiji: “Fiji, Suva” (BMNH).
- flavina* Hreblay et L. Ronkay, 1997 – Described as *Gortyna flavina* in HREBLAY & RONKAY (1997: 66). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 1 km W of Tatachia peak” (CGYF).
- flavitincta* Hreblay et L. Ronkay, 1995 – Described as *Dasypolia (Dasypolia) flavitincta* in HREBLAY & RONKAY (1995b: 358). Holotype male, China: “China, Tibet: 4 km S of Karru-Ochen Tso” (HNHM).
- flavoguttata* Hreblay et L. Ronkay, 1999 – Described as *Himalistra flavoguttata* in HREBLAY & RONKAY (1999: 532). Holotype male, Nepal: “West-Nepal, 20 km N of Dailekh” (CMH).
- flora* Hreblay, 1996 – Described as *Perigrapha (Opacographa) flora* in HREBLAY (1996a: 73). Holotype male, Turkmenistan: “Turkmenistan, Kopet-Dagh Mts, 6 km Ipay-Kala” (CMH).
- forcipata* Hreblay et L. Ronkay, 1998 – Described as *Meganocytycia forcipata* in HREBLAY & RONKAY (1998: 208). Holotype male, Nepal: “Nepal, Ganesh Himal, 12 km S from Somdang” (CMH).
- formontana* Hreblay et L. Ronkay, 1997 – Described as *Hermonassa formontana* in HREBLAY & RONKAY (1997: 22). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 5 km SW of Tayüling” (CMH).
- formosana* Kobayashi et Hreblay, 1998 – Described as *Nyctycia mesomelana formosana* in KOBAYASHI et al. (1998: 263). Holotype male, Taiwan: “Taiwan, Ilan, Szuyuan-Yakou” (NSMT).
- formosana* Hreblay et L. Ronkay, 1998 – Described as *Himachalia formosana* in HREBLAY & RONKAY (1998: 145). Holotype male, Taiwan: “Taiwan, Prov Pingtung, 1 km S of Shihmen” (CMH).
- formosana* Hreblay et L. Ronkay, 1998 – Described as *Amphipyra formosana* in HREBLAY & RONKAY (1998: 274). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 3 km SW of Tsuifeng” (CMH).
- formosanus* Hreblay et L. Ronkay, 1998 – Described as *Dryobotodes formosanus* in HREBLAY & RONKAY (1998: 185). Holotype male, Taiwan: “Taiwan, Prov. Taitung, 2 km N of Tupan” (CMH).
- fortunata* Hreblay et L. Ronkay, 1997 – Described as *Oroplexia fortunata* in HREBLAY & RONKAY (1997: 62). Holotype female, Taiwan: “Taiwan, Prov. Miaoli, 49 km E of Tungshih” (CMH).
- frontieri* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Potnyctycia frontieri* in HREBLAY et al. (1999b: 24). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- fumata* Hreblay et Plante, 1994 – Described as *Himalistra fusca fumata* in HREBLAY et al. (1994: 262). Holotype female, Pakistan: “N-Pakistan, 10 km SW of Astor, Rama” (CMH).

- furiosa* Hreblay et L. Ronkay, 1999 – Described as *Lithophane furiosa* in HREBLAY & RONKAY (1999: 524). Holotype female, Nepal: “West-Nepal, 21 km N of Dailekh” (CMH).
- fusca* Hreblay et Yoshimatsu, 1996 – Described as *Mythimna (Morphopoliana) stolidus fusca* in HREBLAY (1996b: 137). Holotype male, Philippines: “Baguio Mtn. Prov., Luzon, Philip. Is.” (NIAES).
- fusca* Hreblay et Plante, 1996 – Described as *Oroplexia fusca* in HREBLAY & PLANTE (1996: 667). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, 5 km E of Lukla” (CJP).
- fusca* Hreblay et L. Ronkay, 1998 – Described as *Eugnorisma fusca* in HREBLAY & RONKAY (1998: 142). Holotype male, Nepal: “Nepal, Ganesh Himal, near Godlang” (CGR).
- fuscicosta* Hreblay et Plante, 1996 – Described as *Dichagyris fuscicosta* in HREBLAY & PLANTE (1996: 664). Holotype male, Nepal: “Nepal, 6 km SW of Kalinchok peak” (CJP).
- fuscicuprea* Hreblay et L. Ronkay, 1998 – Described as *Trichoridia fuscicuprea* in HREBLAY & RONKAY (1998: 178). Holotype female, Nepal: “Nepal, Annapurna Himal, between Ghorepani and Deorali” (HNHM).
- fuscovestita* Hreblay et L. Ronkay, 1998 – Described as *Orthosia reticulata fuscovestita* in HREBLAY & RONKAY (1998: 157). Holotype male, Taiwan: “Taiwan, Prov Miaoli, 49 km E of Tungshih” (CMH).
- gabori* Hreblay, 1992 – Described as *Conistra gabori* in HREBLAY (1992b: 541). Holotype male, Turkmenistan: “Turkmenia, Kopet-Dagh Mts, 15 km s of Nochur” (CMH).
- gandaki* Hreblay et Plante, 1996 – Described as *Oroplexia pumila gandaki* in HREBLAY & PLANTE (1996: 669). Holotype male, Nepal: “C-Nepal, Kali-Gandaki-Tal, Kalopani-Dhampu” (ZSM).
- ganeshgurungi* Hreblay et L. Ronkay, 1998 – Described as *Altipolia ganeshgurungi* in HREBLAY & RONKAY (1998: 198). Holotype male, Nepal: “Ganesh Himal, Khurpudanda pass, W slope” (CGR).
- ganeshi* Hreblay et L. Ronkay, 1998 – Described as *Harutaeographa ganeshi* in HREBLAY & RONKAY (1998: 154). Holotype male, Nepal: “Nepal, Ganesh Himal, 2 km W Gholjong” (CGR).
- ganeshi* Hreblay, 1998 – Described as *Apamea ganeshi* in HREBLAY & RONKAY (1998: 247). Holotype male, Nepal: “Nepal, Ganesh Himal, 1 km E of Gadrang” (CMH).
- gangtoki* Hreblay et L. Ronkay, 1998 – Described as *Apamea (s.l.) gangtoki* in HREBLAY *et al.* (1998b: 145). Holotype male, India: “Sikkim, Gangtok” (MNHU).
- gansuensis* Hreblay et L. Ronkay, 1998 – Described as *Dichagyris gansuensis* in HREBLAY *et al.* (1998b: 121). Holotype male, China: “China, Gansu, Qilian shan, (Jatjugan), Umg, Jingtie-shan” (HNHM).
- garza* Hreblay et L. Ronkay, 1999 – Described as “*Oroplexia*” *garza* in HREBLAY & RONKAY (1999: 562). Holotype male, Nepal: “Nepal, Mt Kalinchok, 5 km W of Bigu” (CMH).
- gemina* Hreblay et L. Ronkay, 1995 – Described as *Estagrotis gemina* in HREBLAY & RONKAY (1995a: 250). Holotype female, China: “S.E. Tibet, Tsangpo valley, Nyima La” (BMNH).
- gepida* Hreblay, 1996 – Described as *Perigrappa (Perigrappa) i-cinctum gepida* in HREBLAY (1996a: 69). Holotype male, France: “Digne” (NHMW).
- glauca* Hreblay et L. Ronkay, 1998 – Described as *Lithophane glauca* in HREBLAY & RONKAY (1998: 179). Holotype male, Nepal: “Annapurna Himal, 1.5 km SE of Nangethanti” (CGR).
- glaucozona* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Xanthia glaucozona* in HREBLAY *et al.* (1999b: 59). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- glenurina* Hreblay et L. Ronkay, 1999 – Described as *Apamea glenurina* in HREBLAY & RONKAY (1999: 556). Holotype male, Nepal: “Nepal, Arun valley, 12 km N of Chitre” (CMH).
- goliath* Hreblay et Plante, 1998 – Described as *Isochlora goliath* in HREBLAY *et al.* (1998b: 108). Holotype male, China: “China, Tibet, 20 km E of Old Tingri” (CMH).

- goperma* Hreblay et L. Ronkay, 1999 – Described as *Apamea goperma* in HREBLAY & RONKAY (1999: 555). Holotype male, Nepal: “Nepal, 6 km SW of Kalinchok peak” (CMH).
- gorza* Hreblay et L. Ronkay, 1999 – Described as *Agrochola gorza* in HREBLAY & RONKAY (1999: 527). Holotype female, Nepal: “West-Nepal, 11 km N of Dailekh” (CMH).
- grata* Hreblay, 1996 – Described as *Mythimna (Hyphilare) grata* in HREBLAY *et al.* (1996: 99). Holotype male, Nepal: “Nepal, Langtang, 5 km NNE of Dhunche, Barkhu” (CMH).
- gratissima* Hreblay et L. Ronkay, 1999 – Described as *Apamea gratissima* in HREBLAY & RONKAY (1999: 552). Holotype male, Nepal: “Nepal, Ganesh Himal, 3 km SE of Somdang” (CMH).
- grisea* Hreblay et Plante, 1995 – Described as *Transeuplexia grisea* in HREBLAY & PLANTE (1995a: 140). Holotype female, Pakistan: “NO-Pakistan, Kaghan Valley, Battakundi” (CJP).
- grisea* Hreblay et Yoshimatsu, 1996 – Described as *Analetia (Anapoma) grisea* in HREBLAY *et al.* (1996: 122). Holotype male, Nepal: “Nepal, Ganesh Himal, 12 km S from Somdang” (CMH).
- griseana* Hreblay et Plante, 1995 – Described as *Altipolia griseana* in HREBLAY & PLANTE (1995b: 542). Holotype male, Nepal: “Nepal, Taplejung area, Mt Phathibhara” (CMH).
- griseiorbis* Hreblay et L. Ronkay, 1997 – Described as *Cerastis griseiorbis* in HREBLAY & RONKAY (1997: 26). Holotype male, Taiwan: “Taiwan, Prov. Taitung, 2 km E of Hsiangyang” (CGYF).
- griseobrunnea* Hreblay et L. Ronkay, 1999 – Described as *Lithophane griseobrunnea* in HREBLAY & RONKAY (1999: 522). Holotype male, Nepal: “West-Nepal, 20 km N of Dailekh” (CMH).
- grisescens* Hreblay et L. Ronkay, 1998 – Described as *Orthosia grisescens* in HREBLAY & RONKAY (1998: 158). Holotype female, Nepal: “Nepal, Dhaulagiri Himal, 2 km S of Lebang” (CMH).
- hackeri* Hreblay et L. Ronkay, 1995 – Described as *Himalistra hackeri* in HREBLAY *et al.* (1995: 53). Holotype male, Nepal: “Nepal, Langtang, between Cholang Pati and Dimsa” (HNHM).
- hackeri* Hreblay et Yoshimatsu, 1996 – Described as *Mythimna (Mythimna) hackeri* in HREBLAY *et al.* (1996: 90). Holotype male, Nepal: “Nepal, Ganesh Himal, 1 km E of Yurekharka” (CMH).
- hackeri* Hreblay et L. Ronkay, 1998 – Described as *Owadaglaea hackeri* in HREBLAY & RONKAY (1998: 225). Holotype male, Nepal: “Ganesh Himal, above Nesim” (CGR).
- harpegnoma* Hreblay et L. Ronkay, 1998 – Described as *Xestia harpegnoma* in HREBLAY *et al.* (1998b: 127). Holotype male, China: “China, Gansu, Xiahe” (HNHM).
- heidi* Hreblay, 1996 – Described as *Perigrapha (Perigrapha) heidi* in HREBLAY (1996a: 70). Holotype male, Tajikistan: “USSR, Tadzhikistan, Hissarskij Mts, Romit” (ZMH).
- heinickei* Hreblay, 1998 – Described as *Apamea heinickei* in HREBLAY & RONKAY (1998: 246). Holotype male, Nepal: “Nepal, Kalinchok area, 6 km NNE of Muldi (Murre)” (CMH).
- herczigi* Hreblay et L. Ronkay, 1998 – Described as *Amphipyra herczigi* in HREBLAY & RONKAY (1998: 273). Holotype male, Nepal: “Nepal, Langtang, near Chandrabari” (HNHM).
- herczigi* Hreblay et L. Ronkay, 2000 – Described as *Cryphia herczigi* in HREBLAY & RONKAY (2000: 2). Holotype male, Taiwan: “Taiwan, Prov. Taichung, Anmashan, Tashueshan FRA” (HNHM).
- hermanni* Hreblay et L. Ronkay, 1998 – Described as *Elwesia parallela hermanni* in HREBLAY & RONKAY (1998: 235). Holotype male, India: “India, Himachal Pradesh, 15 km W of Shimla” (CGR).
- hiemalis* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Nyctycia endoi hiemalis* in HREBLAY *et al.* (1999b: 16). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- himacola* Hreblay et Legrain, 1999 – Described as *Analetia (Anapoma) himacola* in HREBLAY *et al.* (1999a: 387). Holotype male, Nepal: “West-Nepal, 11 km N of Dailekh” (CMH).
- himalaya* Hreblay et L. Ronkay, 1998 – Described as *Orthosia limbata himalaya* in HREBLAY & RONKAY (1998: 157). Holotype male, Nepal: “Nepal, Ganesh Himal, 1 km N of Nesim” (CMH).

- himalayensis* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Dryobotodes himalayensis* in HREBLAY *et al.* (1999b: 6). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- himalayica* Gyulai et Hreblay, 2001 – Described as *Trichanarta himalayica* in GYULAI & HREBLAY (2001: 710). Holotype male, China: “Tibet West, Himalaya Mts, 20 km West of Namru” (HNHM-PGY).
- hoenei* Hreblay et Kononenko, 1999 – Described as *Diphtherocome hoenei* in HREBLAY & KONONENKO (1999: 671). Holotype male, China: “[China, Prov. Shensi], Tapaishan im Tsiling, Sued-Shenshi” (MAK).
- hohuana* Hreblay et L. Ronkay, 2000 – Described as *Cryphia hohuana* in HREBLAY & RONKAY (2000: 3). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 3 km SW of Tsuifeng” (CMH).
- huberti* Hreblay et L. Ronkay, 1999 – Described as *Orthosia huberti* in HREBLAY & RONKAY (1999: 5498). Holotype male, Vietnam: “Vietnam, Tonkin, Fan-si-pan” (CMH).
- humilis* Hreblay et L. Ronkay, 1998 – Described as *Phlogophora humilis* in HREBLAY & RONKAY (1998: 259). Holotype male, Nepal: “Nepal, Taplejung area, Lal Kharka” (CMH).
- hyphilarioides* Hreblay, 1996 – Described as *Analetia (Anapoma) hyphilarioides* in HREBLAY *et al.* (1996: 122). Holotype male, Nepal: “Nepal, Ganesh Himal, 12 km S from Somdang” (CMH).
- hypobscura* Hreblay et L. Ronkay, 1998 – Described as *Euxoa hypobscura* in HREBLAY *et al.* (1998b: 113). Holotype male, China: “China, Tibet, Tsou La (Pass)” (CJP).
- ignifera* Hreblay, 1998 – Described as *Mythimna (Hyphilare) ignifera* in HREBLAY *et al.* (1998a: 389). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok” (CMH).
- ignorata* Hreblay, 1996 – Described as *Haderonia ignorata* in HREBLAY (1996c: 70). Holotype male, China: “China, Pu-tsu-fang” (BMNH).
- ignorata* Hreblay et Yoshimatsu, 1998 – Described as *Mythimna (Hyphilare) ignorata* in HREBLAY *et al.* (1998a: 392). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok” (CMH).
- ignota* Hreblay, 1998 – Described as *Mythimna (Hyphilare) ignota* in HREBLAY *et al.* (1998a: 391). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok” (CMH).
- illustris* Hreblay, 1993 – Described as *Orthosia (Monima) cruda illustris* in HREBLAY (1993: 78). Holotype male, Turkey: “Türkei, P. Adana, 5 km W of Damlama” (CMH).
- imitabilis* Hreblay, 1993 – Described as *Orthosia (Monima) imitabilis* in HREBLAY (1993: 76). Holotype male, Turkmenistan: “USSR, Türkmenia, Kopet-Dag Mts, 5 km NW of Tutlikala” (CMH).
- imitans* Hreblay et L. Ronkay, 1997 – Described as *Gortyna imitans* in HREBLAY & RONKAY (1997: 80). Holotype male, China: “Li-kiang (China), Provinz Nord-Yuennan” (MAK).
- impar* Hreblay et Plante, 1998 – Described as *Athetis impar* in HREBLAY *et al.* (1998b: 152). Holotype male, China: “China, Tibet, 10 km E of Nyemo” (CJP).
- impectinata* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Diphtherocome impectinata* in HREBLAY *et al.* (1999b: 82). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- implicata* Hreblay et L. Ronkay, 1998 – Described as *Himalistra implicata* in HREBLAY & RONKAY (1998: 196). Holotype female, Nepal: “Ganesh Himal, 7 km of W Godlang” (CGR).
- indistincta* Hreblay et L. Ronkay, 1998 – Described as *Lithopolia indistincta* in HREBLAY & RONKAY (1998: 162). Holotype male, Nepal: “Nepal, Annapurna region, Banthanti” (CMH).
- infenestra* Hreblay et L. Ronkay, 1998 – Described as *Hyalobole infenestra* in HREBLAY & RONKAY (1998: 239). Holotype male, Nepal: “Nepal, Annapurna Himal, 2 km E of Ghorepani” (HNHM).
- inflata* Hreblay et Plante, 1996 – Described as *Oroplexia inflata* in HREBLAY & PLANTE (1996: 666). Holotype male, Nepal: “Nepal, 4 km SW of Kalinchok peak” (CMH).

- informis* Hreblay et L. Ronkay, 1998 – Described as *Mniotype informis* in HREBLAY & RONKAY (1998: 192). Holotype male, Nepal: “Nepal, Arun valley: 11 km N of Hille” (CMH).
- inthanoni* Hreblay et L. Ronkay, 1999 – Described as *Mniotype inthanoni* in HREBLAY & RONKAY (1999: 551). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Inthanon” (CMH).
- intolerabilis* Hreblay, 1992 – Described as *Mythimna (Aletia) intolerabilis* in HREBLAY (1992a: 522). Holotype male, China: “China, N-Sinkiang, Wu-lu-mu-ch’i” (CJP).
- inversa* Hreblay et L. Ronkay, 1999 – Described as *Lithopolia (Thaipolia) inversa* in HREBLAY & RONKAY (1999: 513). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok, 16 km NW of Fang” (CMH).
- ionea* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Hemiglaea ionea* in HREBLAY et al. (1999b: 49). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- ionochlora* L. Ronkay, Varga et Hreblay, 1998 – Described as *Haemerusia ionochlora* in RONKAY et al. (1998: 276). Holotype female, Turkmenistan: “USSR, Turkmenia, Kara-Kum desert, 50 km N of Ashgabat” (CGR).
- ionochlora* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Nyctycia ionochlora* in HREBLAY et al. (1999b: 18). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- ipaykala* L. Ronkay, Varga et Hreblay, 1998 – Described as *Dasytopia ipaykala* in RONKAY et al. (1998: 237). Holotype female, Turkmenistan: “Turkmenistan, Kopet-Dagh Mts, 6 km S of Ipaj-Kala” (CMH).
- isolata* Hreblay et Plante, 1995 – Described as *Xenotrachea isolata* in HREBLAY & PLANTE (1995b: 543). Holotype male, Nepal: “Nepal, Taplejung area, Tambowa” (CJP).
- izabella* Hreblay et L. Ronkay, 1998 – Described as *Harutaographa izabella* in HREBLAY & RONKAY (1998: 153). Holotype male, Nepal: “Nepal, Annapurna Himal, 1 km E of Ghorepani” (CMH).
- jacktordoffi* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Isolasia jacktordoffi* in HREBLAY et al. (1999b: 31). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- javacorna* Hreblay et Legrain, 1998 – Described as *Mythimna (Hyphilare) javacorna* in HREBLAY et al. (1998a: 395). Holotype male, Indonesia: “[Indonesien] W Java, Mt Gedeh” (BMNH).
- jeleni* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Himalistra jeleni* in HREBLAY et al. (1999b: 9). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 6 km W Sa Pa” (HNHM).
- jumla* Hreblay et L. Ronkay, 1999 – Described as *Dasytopia (Dasymixis) jumla* in HREBLAY & RONKAY (1999: 521). Holotype male, Nepal: “West-Nepal, 21 km N of Dailekh” (CMH).
- jumla* Hreblay et L. Ronkay, 1999 – Described as *Hemiglaea jumla* in HREBLAY & RONKAY (1999: 545). Holotype male, Nepal: “West-Nepal, 20 km N of Dailekh” (CMH).
- jutka* Hreblay, 1990 – Described as *Mythimna (Aletia) jutka* in HREBLAY (1990: 125). Holotype male, Mongolia: “Mongolia, Central aimak, Tal des Tola, zwischen Somon Altanbulag und Somon Tariat, 30 km ONO von Tariat” (HNHM).
- jutka* Hreblay et L. Ronkay, 1998 – Described as *Karana jutka* in HREBLAY & RONKAY (1998: 267). Holotype male, Nepal: “Nepal, Taplejung area, Lal Kharka” (CMH).
- kaghanensis* Hreblay et L. Ronkay, 1999 – Described as *Harutaographa bidui kaghanensis* in HREBLAY & RONKAY (1999: 493). Holotype male, Pakistan: “Pakistan, NW-Frontier, Kagan valley, Khanian village” (CMH).

- karma* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Agrochola karma* in HREBLAY *et al.* (1999b: 45). Holotype female, Nepal: “West-Nepal, 20 km N of Dailekh” (CMH).
- karma* Hreblay et L. Ronkay, 1999 – Described as *Nyctycia karma* in HREBLAY & RONKAY (1999: 536). Holotype male, Nepal: “East-Nepal, Deorali Danda, 6 km NW of Yamphudin” (CMH).
- kathmandica* Hreblay, Legrain et Yoshimatsu, 1996 – Described as *Leucania kathmandica* in HREBLAY *et al.* (1996: 124). Holotype male, Nepal: “Nepal, Kathmandu valley, Godavari” (ZSM).
- kautti* Hreblay et L. Ronkay, 1998 – Described as *Euxoa amorpha kautti* in HREBLAY *et al.* (1998b: 114). Holotype male, India: “Indien, Himachal Pradesh, Spiti, Spiti valley, Kaza” (CGR).
- keracola* Hreblay, 1999 – Described as *Analetia (Anapoma) himacola keracola* in HREBLAY *et al.* (1999a: 387). Holotype male, India: “S-India, Kerala, 7 km N Munnar, Eravikulam, NP” (CMH).
- kerala* Hreblay et Legrain, 1999 – Described as *Mythimna (Hyphilare) kerala* in HREBLAY *et al.* (1999a: 381). Holotype male, India: “S-India, Kerala, 7 km N Munnar, Eravikulam, NP” (CMH).
- khangari* Hreblay et L. Ronkay, 1999 – Described as *Dasypolia khangari* in HREBLAY & RONKAY (1999: 519). Holotype female, Nepal: “Nepal, Annapurna Himal, 4 km NW of Khangar” (CMH).
- khasiensis* Hreblay et Legrain, 1998 – Described as *Mythimna (Mythimna) khasiensis* in HREBLAY *et al.* (1998a: 383). Holotype male, India: “Khasia Hills, Assam, Nissary” (BMNH).
- kofka* Hreblay, 1996 – Described as *Perigrapha (Harutaenographa) kofka* in HREBLAY (1996a: 77). Holotype male, India: “Muktesar, Naini Tal” (BMNH).
- kononenkoi* Hreblay et L. Ronkay, 1998 – Described as *Hyalobole kononenkoi* in HREBLAY & RONKAY (1998: 240). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 5 km SW of Tayuling” (CMH).
- kosagezai* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Agrochola kosagezai* in HREBLAY *et al.* (1999b: 39). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- kretania* L. Ronkay, Yela & Hreblay, 2001 – Described as *Polymixis (Myxinia) rufocincta kretania* in RONKAY *et al.* (2001 : 219). Holotype male, Greece: “Crete, 5 km S Hersonissos, 22 km E Heraklion” (CMF).
- krisztina* Hreblay et L. Ronkay, 1998 – Described as *Mniotype krisztina* in HREBLAY *et al.* (1998b: 144). Holotype male, China: “China, Tibet, 20 km S of Saga, 2 km S of Lodu Loney Tso” (CMH).
- kumari* Hreblay et L. Ronkay, 1999 – Described as *Apamea kumari* in HREBLAY & RONKAY (1999: 553). Holotype female, Nepal: “Nepal, Ganesh Himal, 2 km E of Thangjet” (CMH).
- kunandrasi* Hreblay et L. Ronkay, 1999 – Described as *Agrochola kunandrasi* in HREBLAY & RONKAY (1999: 529). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Sa Pa, Fan-si-pan Mts, Frontier Satellite Camp FTC” (HNHM).
- kuni* Hreblay et L. Ronkay, 1999 – Described as *Diphtherocome kuni* in HREBLAY & RONKAY (1999: 565). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- lama* Hreblay et L. Ronkay, 1999 – Described as *Nyctycia lama* in HREBLAY & RONKAY (1999: 534). Holotype female, Nepal: “West-Nepal, 14 km N of Dailekh” (CMH).
- lamprosericea* Hreblay et L. Ronkay, 1998 – Described as *Agrotis lamprosericea* in HREBLAY *et al.* (1998b: 119). Holotype male, China: “Gyantse” (MNHU).
- larga* Hreblay, 1998 – Described as *Mythimna (Hyphilare) larga* in HREBLAY *et al.* (1998a: 400). Holotype male, Indonesia: “[Indonesien] Sumatra, Sitahoan” (BMNH).

- laszlogabi* Hreblay et L. Ronkay, 2000 – Described as *Odontestera laszlogabi* in HREBLAY & RONKAY (2000: 8). Holotype male, Taiwan: “Taiwan, Prov. Ilan, Ming Chyr Forest Recreation Area” (CGR).
- latens* Hreblay et L. Ronkay, 1998 – Described as *Blepharosis latens* in HREBLAY *et al.* (1998b: 140). Holotype male, China: “Li-kiang (China), Provinz Nord-Yuennan” (MAK).
- laudeti* Hreblay et L. Ronkay, 1998 – Described as *Charanyctycia laudeti* in HREBLAY & RONKAY (1998: 218). Holotype male, Nepal: “Nepal, Ganesh Himal, between Godlang and Nesim” (CGR).
- laxa* Hreblay et Yoshimatsu, 1996 – Described as *Mythimna (Hyphilare) laxa* in HREBLAY *et al.* (1996: 107). Holotype male, Nepal: “Nepal, Ganesh Himal, 2 km E of Thangjet” (CMH).
- lepida* Hreblay, Legrain et Yoshimatsu, 1996 – Described as *Mythimna (Hyphilare) lepida* in HREBLAY *et al.* (1996: 108). Holotype male, India: “Khasis” (BMNH).
- leptographa* Hreblay et L. Ronkay, 1995 – Described as *Dasyptolia (Dasyptolia) leptographa* in HREBLAY & RONKAY (1995b: 363). Holotype female, China: “China, Tibet: 4 km S of Karru-Ochen Tso” (CJP).
- leptoaenia* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Hemiglaea costalis leptoaenia* in HREBLAY *et al.* (1999b: 48). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- leta* Hreblay et L. Ronkay, 1999 – Described as *Dasyptolia leta* in HREBLAY & RONKAY (1999: 517). Holotype female, Nepal: “Nepal, Annapurna Himal, 4 km SE of Kaisang” (CMH).
- leucocollaris* Hreblay et L. Ronkay, 1998 – Described as *Rhynchaglaea hemixantha leucocollaris* in HREBLAY & RONKAY (1998: 223). Holotype female, Nepal: “Ganesh Himal, near Nesim” (CGR).
- lieni* Hreblay, 1998 – Described as *Apamea lieni* in HREBLAY & RONKAY (1998: 247). Holotype male, Taiwan: “Taiwan, Prov. Taoyuan, 16 km E of Fuhsing” (CMH).
- likianga* Hreblay et Kononenko, 1999 – Described as *Diphtherocome hoenei likianga* in HREBLAY & KONONENKO (1999: 672). Holotype male, China: “[China, Prov. Yunnan], Likiang, Prov. Nord-Yuennan” (MAK).
- lilacina* Hreblay et L. Ronkay, 1998 – Described as *Euplexia lilacina* in HREBLAY & RONKAY (1998: 261). Holotype male, Nepal: “Ganesh Himal, near Yurekharka” (HNHM).
- lineolata* Hreblay et L. Ronkay, 1998 – Described as *Eugnorisma lineolata* in HREBLAY & RONKAY (1998: 140). Holotype male, Nepal: “Nepal, Langtang, near Chandrabari” (CGR).
- lithoplana* Hreblay et L. Ronkay, 1998 – Described as *Xestia (s.l.) lithoplana* in HREBLAY *et al.* (1998b: 128). Holotype male, China: “Chumbitang” (MNHU).
- longipennis* Hreblay et L. Ronkay, 1998 – Described as *Hemiglaea longipennis* in HREBLAY & RONKAY (1998: 221). Holotype male, Nepal: “Ganesh Himal, near Slya” (CGR).
- longirostris* Hreblay et L. Ronkay, 1998 – Described as *Hyalobole changae longirostris* in HREBLAY & RONKAY (1998: 242). Holotype female, Nepal: “Nepal, Ganesh Himal, near Nesim” (CGR).
- longivalva* Hreblay et Kononenko, 1999 – Described as *Diphtherocome longivalva* in HREBLAY & KONONENKO (1999: 677). Holotype male, China: “[China, Prov. Yunnan], Likiang, Prov. Nord-Yuennan” (MAK).
- lucida* Yoshimatsu et Hreblay, 1996 – Described as *Mythimna (Mythimna) lucida* in YOSHIMATSU & HREBLAY (1996b: 265). Holotype male, Taiwan: “Chun Yang, Nantou Hsien, Taiwan” (NIAES).
- lucida* Hreblay et L. Ronkay, 1998 – Described as *Owadaglaea lucida* in HREBLAY & RONKAY (1998: 229). Holotype male, Nepal: “Nepal, Arun valley, 22 km N of Hille” (CMH).
- luna* Hreblay et L. Ronkay, 1999 – Described as *Trichoridia luna* in HREBLAY & RONKAY (1999: 516). Holotype male, Nepal: “Nepal, Ganesh Himal, Khurpudanda Pass” (CMH).

- lupsangwangi* Hreblay et L. Ronkay, 1998 – Described as *Agrotis lamprosericea lupsangwangi* in HREBLAY *et al.* (1998b: 119). Holotype male, China: “China, Tibet, 1 km S of Paiku Tso” (CJP).
- luteomixta* Hreblay et L. Ronkay, 1998 – Described as *Rhynchaglaea luteomixta* in HREBLAY & RONKAY (1998: 223). Holotype male, Taiwan: “Taiwan, Prov. Taoyuan, Ming Chyr Forest Recreation Area” (CGYF).
- machka* Hreblay et L. Ronkay, 1998 – Described as *Blepharomorpha machka* in HREBLAY *et al.* (1998b: 143). Holotype male, Pakistan: “Pakistan, Himalaya Mts, Kaghan valley, Tathabaya” (HNHM).
- macilenta* Hreblay et L. Ronkay, 1998 – Described as *Orthosia macilenta* in HREBLAY & RONKAY (1998: 158). Holotype male, Nepal: “Nepal, Ganesh Himal, 1 km E of Gadrang” (CMH).
- macroscura* Hreblay et L. Ronkay, 1998 – Described as *Agrotis macroscura* in HREBLAY *et al.* (1998b: 116). Holotype male, China: “China, Tibet, 10 km E of Nyemo” (CJP).
- macrophthalmalma* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Elwesia macrophthalmalma* in HREBLAY *et al.* (1999b: 54). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- major* Hreblay et Thöny, 1995 – Described as *Checupa major* in HREBLAY & THÖNY (1995: 24). Holotype male, Philippines: “Philippines, Luzon, Camarines sud, Naga Relay Stn, Mount Isarog, Ocampo” (CWS).
- maminti* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Chandata maminti* in HREBLAY *et al.* (1999b: 78). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 3 km NW Cat Cat” (HNHM).
- manfredi* Hreblay, 1994 – Described as *Orthosia (Orthosia) manfredi* in HREBLAY (1994: 242). Holotype male, Morocco: “Morocco, Middle-Atlas Mts, 6 km SE Azrou” (CMH).
- manfredi* Hreblay et L. Ronkay, 1998 – Described as *Auchmis manfredi* in HREBLAY & RONKAY (1998: 251). Holotype male, Nepal: “Nepal, Annapurna Himal, 11 km SW of Jomsom, Noma pasture” (HNHM).
- manopi* Hreblay, 1998 – Described as *Mythimna (Hyphilare) manopi* in HREBLAY *et al.* (1998a: 392). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok” (CMH).
- marcida* Hreblay et L. Ronkay, 2000 – Described as *Karana hoenei marcida* in HREBLAY & RONKAY (2000: 33). Holotype male, Taiwan: “Taiwan, Prov. Taoyuan, 14 km E of Fuhshing” (CMH).
- marginalis* Hreblay et L. Ronkay, 1998 – Described as *Hyalobole marginalis* in HREBLAY & RONKAY (1998: 238). Holotype male, Nepal: “Nepal, 6 km NNE of Muldi (Murre)” (CMH).
- marginata* Hreblay et L. Ronkay, 1998 – Described as *Hermonassa marginata* in HREBLAY & RONKAY (1998: 131). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, 14 km E of Lukla” (CMH).
- maria* Hreblay et L. Ronkay, 1999 – Described as *Harutaegerapha maria* in HREBLAY & RONKAY (1999: 494). Holotype male, Pakistan: “Pakistan, Prov. Jammu & Kaschmir, Naltar valley, 5 km E of Naltar village” (CMH).
- marki* Hreblay et L. Ronkay, 2000 – Described as *Fabiania marki* in HREBLAY & RONKAY (2000: 29). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- marmorata* Hreblay et L. Ronkay, 1998 – Described as *Daseuplexia marmorata* in HREBLAY & RONKAY (1998: 209). Holotype female, Nepal: “Annapurna Himal, Ghorepani” (CGR).
- marpha* Hreblay et L. Ronkay, 1999 – Described as *Harutaegerapha marpha* in HREBLAY & RONKAY (1999: 495). Holotype female, Nepal: “Nepal, Dhaulagiri Himal, 6 km NW of Marpha” (CMH).

- matura* Hreblay et L. Ronkay, 1998 – Described as *Oroplexia matura* in HREBLAY *et al.* (1998b: 150). Holotype male, India: “Tsomgo” (MNHU).
- mauritiusi* Hreblay, 1998 – Described as *Mythimna (Morphopoliana) mauritiusi* in HREBLAY *et al.* (1998a: 411). Holotype male, Mauritius: “Curepipe, Mauritius” (BMNH).
- maxima* Hreblay, 1998 – Described as *Mythimna (Hyphilara) maxima* in HREBLAY *et al.* (1998a: 404). Holotype male, Philippines: “Philippinen, Mindanao, S-Mt Kitanglad” (CMH).
- mediofusca* Hreblay et L. Ronkay, 1999 – Described as *Orthosia mediofusca* in HREBLAY & RONKAY (1999: 496). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok, 18 km NW of Fang” (CMH).
- medionigra* Hreblay et L. Ronkay, 1998 – Described as *Parabole medionigra* in HREBLAY & RONKAY (1998: 217). Holotype male, Nepal: “Nepal, Arun valley, 12 km N of Chitre” (CMH).
- megasripta* Hreblay et L. Ronkay, 1998 – Described as *Rhynchaglaea megasripta* in HREBLAY & RONKAY (1998: 222). Holotype male, Nepal: “Ganesh Himal, 2 km W of Gholjong” (CGR).
- meiseri* Hreblay et Thöny, 1995 – Described as *Checupa meiseri* in HREBLAY & THÖNY (1995: 26). Holotype male, Philippines: “Philippinen, Mindanao, S-Mt Kitanglad” (CMH).
- melachlora* Hreblay et L. Ronkay, 1999 – Described as *Euplexidia melachlora* in HREBLAY & RONKAY (1999: 559). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Sa Pa, Fan-si-pan Mts, Frontier Satellite Camp FTC” (HNHM).
- melanura* Hreblay et L. Ronkay, 1999 – Described as *Pyrria umbra melanura* in HREBLAY & RONKAY (1999: 488). Holotype male, Nepal: “Nepal, Annapurna Himal, 1 km E of Khangar” (CMH).
- melanura* Hreblay et L. Ronkay, 1999 – Described as *Agrochola melanura* in HREBLAY & RONKAY (1999: 530). Holotype male, Nepal: “Nepal, Annapurna Himal, 2 km SW of Kaisang” (CMH).
- merkli* Hreblay et L. Ronkay, 1999 – Described as *Kisegira merkli* in HREBLAY & RONKAY (1999: 508). Holotype female, Laos: “LAOS, Prov. Champassak, Dong Hua Xao NBCA, Nam Phak River” (HNHM).
- metallica* Hreblay et L. Ronkay, 1998 – Described as “*Conistra*” *metallica* in HREBLAY & RONKAY (1998: 232). Holotype female, Nepal: “Nepal, Langtang, near Chandrabari” (CGR).
- metexotica* Hreblay et L. Ronkay, 1998 – Described as *Euplexidia metexotica* in HREBLAY & RONKAY (1998: 264). Holotype female, Nepal: “Annapurna Himal, between Ghorepani and Nangethanti” (CGR).
- miaoli* Hreblay et Kobayashi, 1997 – Described as *Allophyes miaoli* in HREBLAY & RONKAY (1997: 42). Holotype female, Taiwan: “Taiwan, Prov Miaoli, 49 km E of Tungshih” (CMH).
- microlitha* Hreblay et L. Ronkay, 1998 – Described as *Amphipyra microlitha* in HREBLAY & RONKAY (1998: 272). Holotype male, Nepal: “Nepal, Lapchi Kang Range, 4 km SW of Tselaphu (Doupseyding)” (CMH).
- mikkolai* Hreblay et L. Ronkay, 1998 – Described as *Apamea mikkolai* in HREBLAY & RONKAY (1998: 244). Holotype male, Nepal: “Ganesh Himal, 2 km E of Thangjet” (CMH).
- minima* Hreblay et L. Ronkay, 1998 – Described as *Isochlora minima* in HREBLAY *et al.* (1998b: 108). Holotype male, Mongolia: “Mongolia, Uvs aimak, 4 km OSO vom Pass Ulan davaa zw. See Örög nuur und Ulangom” (HNHM).
- minima* Hreblay et L. Ronkay, 1999 – Described as *Paranycyctia minima* in HREBLAY & RONKAY (1999: 544). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, 20 km NW of Sop Kha, 2 km S of Kop Dong” (CMH).
- minorata* Hreblay et L. Ronkay, 1999: described as *Agrochola*’ *minorata* in HREBLAY & RONKAY (1999: 530). Holotype male, China: “Lingping, Südchina” (MAK).
- minuta* Hreblay et Plante, 1998 – Described as *Dichagyris minuta* in HREBLAY *et al.* (1998b: 122). Holotype male, China: “China, Tibet, 5 km W of Paiku Tso” (CMH).

- minutoides* L. Ronkay, Varga et Hreblay, 1998 – Described as *Episema minutoides* in RONKAY *et al.* (1998: 238). Holotype male, Turkmenistan: “USSR, Turkmenia, Kopet-Dagh Mts, Kurkuklab, 6 km W Germob” (HNHM).
- mirabilis* Hreblay et L. Ronkay, 1998 – Described as *Hemiglaea mirabilis* in HREBLAY & RONKAY (1998: 219). Holotype male, Nepal: “Ganesh Himal, near Godlang” (CGR).
- miraculosa* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Paranyctycia miraculosa* in HREBLAY *et al.* (1999b: 37). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 3 km NW Cat Cat” (HNHM).
- mixta* Hreblay et Kononenko, 1999 – Described as *Diphtherocome mixta* in HREBLAY & KONONENKO (1999: 674). Holotype male, China: “[China, Prov. Yunnan], Likiang, Prov. Nord-Yuennan” (MAK).
- mixtazona* Hreblay et L. Ronkay, 1992 – Described as *Oncocnemis mixtazona* in GYULAI *et al.* (1992: 40). Holotype male, Afghanistan: “NO-Afghanistan, Wakhan-Tal, Darrah-e-Shaur” (CCN).
- monogramma* Hreblay et L. Ronkay, 1995 – Described as *Dasypolia (Dasypolia) monogramma* in HREBLAY & RONKAY (1995b: 362). Holotype male, China: “China, Tibet, 1 km S of Paiku Tso” (HNHM).
- montigena* Hreblay et Legrain, 1996 – Described as *Mythimna (Morphopoliana) montigena* in HREBLAY (1996b: 147). Holotype male, Tanzania: “Tanganjika, Kilimandjaro, Marangu” (ZSM).
- moriutii* Yoshimatsu et Hreblay, 1996 – Described as *Mythimna (Hyphilare) moriutii* in YOSHIMATSU & HREBLAY (1996a: 13). Holotype male, Thailand: “Doi Inthanon, Chiang Mai, Thailand” (UO).
- munnnari* Hreblay, 1999 – Described as *Mythimna (Hyphilare) munnnari* in HREBLAY *et al.* (1999a: 383). Holotype male, India: “S-India, Kerala, 7 km N Munnar, Eravikulam, NP” (CMH).
- nani* Hreblay et L. Ronkay, 1999 – Described as *Kisegira nani* in HREBLAY & RONKAY (1999: 509). Holotype male, Thailand: “Thailand, Changwat Nan, 30 km E of Pua” (CMH).
- naumanni* Yoshimatsu et Hreblay, 1998 – Described as *Mythimna (Mythimna) naumanni* in YOSHIMATSU & HREBLAY (1998: 11). Holotype male, China: “Li-kiang (China), Provinz Nord-Yuennan” (MAK).
- nebulosa* L. Ronkay, Varga et Hreblay, 1998 – Described as *Dasypolia nebulosa* in RONKAY *et al.* (1998: 234). Holotype male, Turkmenistan: “USSR, Turkmenia, Kopet-Dagh Mts, Firyuza” (HNHM).
- nepalensis* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Amphipyra owadai nepalensis* in HREBLAY *et al.* (1999b: 93). Holotype male, Nepal: “West-Nepal, 11 km N of Dailekh” (CMH).
- nepalichlora* Hreblay et L. Ronkay, 1999 – Described as *Diphtherocome nepalichlora* in HREBLAY & RONKAY (1999: 570). Holotype male, Nepal: “Nepal, Arun valley, 12 km N of Hille” (CMH).
- nepalicola* Hreblay et L. Ronkay, 1998 – Described as *Diarsia hoenei nepalicola* in HREBLAY & RONKAY (1998: 143). Holotype male, Nepal: “Nepal, Lapchi Kang Range, 3 km SW of Tselaphu (Kalow)” (CMH).
- nepalina* Hreblay et Plante, 1995 – Described as *Euchalcia nepalina* in HREBLAY & PLANTE (1995b: 545). Holotype male, Nepal: “Nepal, Ganesh Himal, 1 km S of Somdang” (CMH).
- nepalina* Hreblay et Yoshimatsu, 1996 – Described as *Mythimna (Hyphilare) nepalina* in HREBLAY *et al.* (1996: 102). Holotype male, Nepal: “Nepal, Ganesh Himal, 2 km E of Thangjet” (CMH).
- nepalina* Hreblay et L. Ronkay, 1999 – Described as *Pyrrhia umbra nepalina* in HREBLAY & RONKAY (1999: 487). Holotype male, Nepal: “Nepal, 4 km SW of Kalinchok peak” (CMH).
- nigriclava* Hreblay et L. Ronkay, 1999 – Described as *Nyctycia nigriclava* in HREBLAY & RONKAY (1999: 535). Holotype male, Nepal: “West-Nepal, 20 km N of Dailekh” (CMH).

- nigriclavata* Hreblay et L. Ronkay, 1998 – Described as *Agrotis nigriclavata* in HREBLAY *et al.* (1998b: 117). Holotype male, China: “Samada” (MNHU).
- nigricomma* Hreblay et L. Ronkay, 1998 – Described as *Owadaglaea nigricomma* in HREBLAY & RONKAY (1998: 228). Holotype male, Nepal: “Ganesh Himal, 1 km E of Gadrang” (CMH).
- nigristriga* Hreblay, Legrain et Yoshimatsu, 1998 – Described as *Leucania nigristriga* in HREBLAY *et al.* (1998a: 412). Holotype male, Thailand: “Thailand, Changwat Nan, 25 km N of Bo Luang” (CMH).
- nigrocincta* Hreblay et L. Ronkay, 1997 – Described as *Perigrapha nigrocincta* in HREBLAY & RONKAY (1997: 35). Holotype male, Taiwan: “Taiwan, Prov. Nantou, Meimu” (CGYF).
- nigrogrisea* Hreblay et L. Ronkay, 1998 – Described as *Blepharita nigrogrisea* in HREBLAY & RONKAY (1998: 190). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 5 km SW of Tayuling” (CMH).
- nigromaculata* Hreblay et L. Ronkay, 1998 – Described as *Rhynchaglaea nigromaculata* in HREBLAY & RONKAY (1998: 221). Holotype male, Nepal: “Nepal, Ganesh Himal, near Slya” (CGR).
- nigrostromia* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Apamea nigrostromia* in HREBLAY *et al.* (1999b: 77). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- nivalis* Hreblay et L. Ronkay, 1995 – Described as *Dasypolia* (*s. l.*) *nivalis* in HREBLAY & RONKAY (1995b: 376). Holotype male, China: “China, Tibet: 4 km S of Karru-Ochen Tso” (HNHM).
- nivea* Hreblay et Plante, 1994 – Described as *Himalistra nivea* in HREBLAY *et al.* (1994: 260). Holotype female, Pakistan: “N-Pakistan, 10 km SW of Astor, Rama” (CJP).
- niveifera* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Nyctycia niveifera* in HREBLAY *et al.* (1999b: 23). Holotype female, Thailand: “Thailand, Changwat Nan, 25 km N of Bo Luang” (CMH).
- nivescens* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Nyctycia nivescens* in HREBLAY *et al.* (1999b: 14). Holotype male, Vietnam: “Vietnam, Prov. Vinh Phu, Tam Dao” (HNHM).
- nobilis* Hreblay et L. Ronkay, 1998 – Described as *Phlogophora nobilis* in HREBLAY & RONKAY (1998: 259). Holotype male, Nepal: “Annapurna Himal, Nangethanti” (CGR).
- nyemonis* Hreblay et L. Ronkay, 1998 – Described as *Parexarnis nyemonis* in HREBLAY *et al.* (1998b: 123). Holotype male, China: “China, Tibet, 10 km E of Nyemo” (CMH).
- obfuscata* Hreblay et L. Ronkay, 1998 – Described as *Apamea lateritia obfuscata* in HREBLAY & RONKAY (1998: 244). Holotype male, Nepal: “Annapurna Himal, 5 km SE of Jomsom, Thadung valley” (CGR).
- obscura* Hreblay et Plante, 1995 – Described as *Himalistra obscura* in HREBLAY *et al.* (1995: 50). Holotype male, India: “Indien, Himachal Pradesh, Rohtang-Paß-S-Seite, ob Gulaba” (CMH).
- obscurographa* Hreblay, 1999 – Described as *Mythimna* (*Hyphilare*) *obscurographa* in HREBLAY *et al.* (1999a: 380). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, 2 km S of Kop Dong” (CMH).
- obsoleta* Hreblay et L. Ronkay, 1995 – Described as *Dasypolia* (*Dasypolia*) *obsoleta* in HREBLAY & RONKAY (1995b: 353). Holotype male, China: “China, Tibet: 4 km S of Karru-Ochen Tso” (CJP).
- obsoleta* Hreblay et L. Ronkay, 1998 – Described as *Potnyctycia obsoleta* in HREBLAY & RONKAY (1998: 206). Holotype male, Nepal: “Nepal, Ganesh Himal, Gothen village” (CGR).
- opposita* Hreblay et L. Ronkay, 1998 – Described as *Bornolis opposita* in HREBLAY & RONKAY (1998: 251). Holotype male, Nepal: “Nepal, Taplejung area, Shimbu (Pakora)” (CMH).
- opulenta* Hreblay et L. Ronkay, 1998 – Described as *Auchmis opulenta* in HREBLAY & RONKAY (1998: 253). Holotype female, Nepal: “Nepal, Annapurna Himal, 10 km SE of Jomsom” (CGR).

- orbiculosa* Hreblay et L. Ronkay, 1998 – Described as *Paranyctycia orbiculosa* in HREBLAY & RONKAY (1998: 215). Holotype male, Nepal: “Ganesh Himal, 2 km W of Gholjong” (CGR).
- orias* Hreblay, 1996 – Described as *Perigrapha (Harutaeographa) orias* in HREBLAY (1996a: 75). Holotype female, India: “Darjeeling” (BMNH).
- ornata* Hreblay et L. Ronkay, 1999 – Described as *Egira ornata* in HREBLAY & RONKAY (1999: 501). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Inthanon” (CMH).
- ornata* Hreblay et L. Ronkay, 1999 – Described as *Euplexidia ornata* in HREBLAY & RONKAY (1999: 561). Holotype female, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok, 19 km NW of Fang” (CMH).
- orogena* Hreblay et L. Ronkay, 1995 – Described as *Dasypolia (Dasymixis) orogena* in HREBLAY & RONKAY (1995b: 374). Holotype female, Nepal: “Nepal, Ganesh Himal, 1 km SE from Somdang” (HNHM).
- orophila* Hreblay et L. Ronkay, 1998 – Described as *Cucullia duplicata orophila* in HREBLAY et al. (1998b: 133). Holotype male, China: “China, Tibet, 20 km S of Saga, 2 km S of Lodu Loney Tso” (CMH).
- ossicolor* Hreblay et Plante, 1998 – Described as *Isochlora ossicolor* in HREBLAY et al. (1998b: 109). Holotype male, China: “China, Tibet, Mayum La (Pass), E side” (CJP).
- owadai* Hreblay et L. Ronkay, 1998 – Described as *Dasypolia owadai* in HREBLAY & RONKAY (1998: 185). Holotype male, Nepal: “C. Nepal, Gandaki, Parbat Distr., Modi Khola” (NSMT).
- owadai* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Amphipyra owadai* in HREBLAY et al. (1999b: 92). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- pali* Hreblay et L. Ronkay, 1998 – Described as *Euplexia pali* in HREBLAY & RONKAY (1998: 260). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 3 km SW of Tsuifeng” (CMH).
- pallida* Hreblay et L. Ronkay, 1999 – Described as *Kisegira pallida* in HREBLAY & RONKAY (1999: 509). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, 7 km W of Pa Pae” (CMH).
- pallidilinea* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Agrochola pallidilinea* in HREBLAY et al. (1999b: 42). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- pallidipennis* Hreblay et L. Ronkay, 1998 – Described as *Amphipyra pallidipennis* in HREBLAY & RONKAY (1998: 274). Holotype male, Nepal: “Ganesh Himal, near Godlang” (CGR).
- pallidipennis* Hreblay et L. Ronkay, 1999 – Described as *Diphtherocome pallidipennis* in HREBLAY & RONKAY (1999: 567). Holotype male, Nepal: “Nepal E., Chagam” (HNHM).
- pallidula* Hreblay et Plante, 1998 – Described as *Diphtherocome pallidula* in HREBLAY et al. (1998b: 111). Holotype male, China: “China, Tibet, 10 km E of Nyemo” (CMH).
- pamiricola* Hreblay et Kononenko, 1995 – Described as *Perigrapha pamiricola* in HREBLAY & KONONENKO (1995: 117). Holotype male, Tajikistan: “Tadjikistan, Pamir, Chorong, Hortus Botan” (ZIN).
- papae* Hreblay et L. Ronkay, 1999 – Described as *Egira papae* in HREBLAY & RONKAY (1999: 502). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, 7 km W of Pa Pae” (CMH).
- parallela* Hreblay et L. Ronkay, 1998 – Described as *Elwesia parallela* in HREBLAY & RONKAY (1998: 234). Holotype female, Nepal: “Ganesh Himal, 2 km W of Gholjong” (CGR).
- parashyu* Hreblay et L. Ronkay, 1998 – Described as *Eupsilia parashyu* in HREBLAY & RONKAY (1998: 182). Holotype female, Nepal: “Ganesh Himal, 7 km of W Godlang” (CGR).
- pastellina* Hreblay et Legrain, 1996 – Described as *Mythimna (Mythimna) pastellina* in HREBLAY et al. (1996: 94). Holotype male, India: “Darjiling” (BMNH).

- pectinicornis* Hreblay et L. Ronkay, 1998 – Described as *Xylena tatajiana pectinicornis* in HREBLAY & RONKAY (1998: 182). Holotype female, Nepal: “Nepal, Tanahoun distr., Dhumre, Bimal Nager” (CMH).
- pectinosa* Hreblay et Plante, 1996 – Described as *Oroplexia pectinosa* in HREBLAY & PLANTE (1996: 669). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, Lukla” (CMH).
- penicillata* Hreblay et L. Ronkay, 1999 – Described as *Kisegira penicillata* in HREBLAY & RONKAY (1999: 505). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, 20 km NW of Mae Ai” (CMH).
- peregovitsi* Hreblay et L. Ronkay, 1999 – Described as *Orthosia peregovitsi* in HREBLAY & RONKAY (1999: 498). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 6 km W Cat Cat” (HNHM).
- pericaspicus* L. Ronkay, Varga et Hreblay, 1998 – Described as *Polymixis pericaspicus* in RONKAY *et al.* (1998: 239). Holotype male, Turkmenistan: “USSR, Turkmenia, Kopet-Dagh Mts, Kurkulab, 6 km W Germob” (HNHM).
- periculosa* Hreblay et L. Ronkay, 1998 – Described as *Euplexidia periculosa* in HREBLAY & RONKAY (1998: 265). Holotype male, India: “Darjiling” (MNHU).
- perigraphoides* Hreblay et L. Ronkay, 1999 – Described as *Egira perigraphoides* in HREBLAY & RONKAY (1999: 500). Holotype male, Thailand: “Thailand, Changwat Mae Hong Song, 21 km NW of Pai” (CMH).
- perseverans* Hreblay et L. Ronkay, 1998 – Described as *Dasyrgeres perseverans* in HREBLAY & RONKAY (1998: 176). Holotype male, Nepal: “Annapurna Himal, between Ghorepani and Deorali” (CGR).
- petra* Hreblay et Yoshimatsu, 1999 – Described as *Leucania petra* in HREBLAY *et al.* (1999a: 384). Holotype male, India: “S-India, Kerala, 7 km N Munnar, Eravikulam, NP” (CMH).
- phahompoki* Hreblay et L. Ronkay, 1999 – Described as *Egira phahompoki* in HREBLAY & RONKAY (1999: 503). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok, 18 km NW of Fang” (CMH).
- picurka* Hreblay et L. Ronkay, 1995 – Described as *Dasypolia (s. l.) picurka* in HREBLAY & RONKAY (1995b: 371). Holotype female, Nepal: “Nepal, Annapurna region, 1 km E from Gorepani” (HNHM).
- pinkeri* Hreblay et Varga, 1993 – Described as *Dioszeghyana schmidtii pinkeri* in HREBLAY (1993: 81). Holotype female, Turkey: “Türkei, P. Ankara. Dütözü Köyü” (CMH).
- pinkisherpani* Hreblay et L. Ronkay, 1998 – Described as *Harutaeographa pinkisherpani* in HREBLAY & RONKAY (1998: 154). Holotype male, Nepal: “Nepal, Ganesh Himal, 2 km SW of Haku” (CGR).
- plantei* Hreblay et Yoshimatsu, 1996 – Described as *Mythimna (Morphopoliana) plantei* in HREBLAY (1996b: 144). Holotype female, Taiwan: “Taiwan, Li-Shan, (Taichung Cy)” (CJP).
- plantei* Hreblay et L. Ronkay, 1997 – Described as *Hermonassa plantei* in HREBLAY & RONKAY (1997: 24). Holotype male, Taiwan: “Taiwan, Prov. Taitung, Hsiangyang” (CMH).
- plumbeata* Hreblay et L. Ronkay, 1997 – Described as *Gortyna plumbeata* in HREBLAY & RONKAY (1997: 78). Holotype male, Thailand: “N-Thailand, Nan Prov., Pua Doi Phu Kha, km 35” (MAK).
- plumbeata* Hreblay et L. Ronkay, 1998 – Described as *Perigrapha (Anorthoa) munda plumbeata* in HREBLAY & RONKAY (1998: 152). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 1 km W of Tatachia peak” (CMH).
- plumbeopaca* Hreblay et L. Ronkay, 2000 – Described as *Xylena plumbeopaca* in HREBLAY & RONKAY (2000: 17). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 3 km SW of Tsuifeng” (CMH).

- plumbitincta* Hreblay et L. Ronkay, 1997 – Described as *Gortyna plumbitincta* in HREBLAY & RONKAY (1997: 77). Holotype male, Taiwan: “Taiwan, Prov. Taitung, Hsiangyang, police station” (CGYF).
- plumbitincta* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Agrochola plumbitincta* in HREBLAY *et al.* (1999b: 46). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- poecila* Hreblay et L. Ronkay, 1997 – Described as *Cosmia poecila* in HREBLAY & RONKAY (1997: 63). Holotype male, Taiwan: “Taiwan, Prov. Taitung, 2 km N of Tupan” (CMH).
- porphyrea* Hreblay et L. Ronkay, 1998 – Described as *Potnyctycia porphyrea* in HREBLAY & RONKAY (1998: 205). Holotype male, Nepal: “Annapurna Himal, Ghorepani” (CGR).
- porphyrea* Hreblay et L. Ronkay, 1998 – Described as *Amphipyra porphyrea* in HREBLAY & RONKAY (1998: 271). Holotype male, Nepal: “Annapurna Himal, 11 km SE of Jomsom, Noma pasture” (CGR).
- porphyrea* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Daseutype porphyrea* in HREBLAY *et al.* (1999b: 37). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 3 km NW Cat Cat” (HNHM).
- prima* Hreblay et L. Ronkay, 1998 – Described as *Karana prima* in HREBLAY & RONKAY (1998: 268). Holotype male, China: “Siao-Lou” (BMNH).
- pseudodisca* Hreblay et L. Ronkay, 1998 – Described as *Xenotrachea albidisca pseudodisca* in HREBLAY & RONKAY (1998: 269). Holotype male, Taiwan: “Taiwan, Prov. Miaoli, 20 km E of Tungshih” (CMH).
- pseudopartita* Hreblay et L. Ronkay, 1998 – Described as *Chandata pseudopartita* in HREBLAY & RONKAY (1998: 263). Holotype male, Nepal: “Nepal, Ganesh Himal, 2 km E of Thangjet” (CMH).
- pseudovivida* Hreblay et L. Ronkay, 1999 – Described as *Diphtherocombe pseudovivida* in HREBLAY & RONKAY (1999: 568). Holotype male, Nepal: “East-Nepal, Deorali Danda, Torontan” (CMH).
- pua* Hreblay et L. Ronkay, 1999 – Described as *Lithopolia (Hiemopolia) pua* in HREBLAY & RONKAY (1999: 512). Holotype male, Thailand: “Thailand, Changwat Nan, 30 km E of Pua” (CMH).
- pulla* Hreblay et L. Ronkay, 2000 – Described as *Fabiania pulla* in HREBLAY & RONKAY (2000: 25). Holotype male, Taiwan: “Taiwan, Prov. Ilan, Ming Chyr Forest Recreation Area” (CGYF).
- pumila* Hreblay et Plante, 1996 – Described as *Oroplexia pumila* in HREBLAY & PLANTE (1996: 668). Holotype male, Nepal: “Nepal, Solu Khumbu Himal: 3 [k]m, Lukla” (CJP).
- punctilinea* Hreblay et L. Ronkay, 1999 – Described as *Agrochola punctilinea* in HREBLAY & RONKAY (1999: 525). Holotype female, Nepal: “East-Nepal, Deorali Danda, Dhupi Pass” (CMH).
- pygmaea* Hreblay et L. Ronkay, 1995 – Described as *Dasytopia (Dasythorax) pygmaea* in HREBLAY & RONKAY (1995b: 375). Holotype female, Pakistan: “Pakistan, Shandur Pass” (CJP).
- pygmaea* Hreblay et L. Ronkay, 1999 – Described as *Nyctycia stenoptera pygmaea* in HREBLAY & RONKAY (1999: 536). Holotype male, Nepal: “East-Nepal, Surke Danda, 1 km W of Kesawa” (CMH).
- quasimodo* Hreblay, 1998 – Described as *Mythimna (Hyphilare) quasimodo* in HREBLAY *et al.* (1998a: 400). Holotype male, Indonesia: “[Indonesien] Sumatra, Aceh” (CHH).
- radiata* Hreblay et L. Ronkay, 2000 – Described as *Hemiglaea radiata* in HREBLAY & RONKAY (2000: 21). Holotype female, Taiwan: “Taiwan, Prov. Tai-Tung, Yakou” (CGYF).
- rai* Hreblay et L. Ronkay, 1995 – Described as *Estagrotis benescripta rai* in HREBLAY & RONKAY (1995a: 245). Holotype male, Nepal: “Nepal, Taplejung area, Mt Phathibhara” (HNHM).
- rama* Hreblay et Plante, 1996 – Described as *Perigrapha (Harutaographa) rama* in HREBLAY (1996a: 78). Holotype male, Pakistan: “N-Pakistan, 10 km SW of Astor, Rama” (CMH).

- ratna* Hreblay et L. Ronkay, 1998 – Described as *Euxoa ratna* in HREBLAY *et al.* (1998b: 114). Holotype male, Nepal: “Nepal, Annapurna Himal, Mesokantu Pass” (CGR).
- rectificata* Hreblay et Plante, 1995 – Described as *Apamea rectificata* in HREBLAY & PLANTE (1995b: 544). Holotype male, Nepal: “Nepal, Langtang, Kyangjin Gompa” (CJP).
- rectilinea* Hreblay et L. Ronkay, 1998 – Described as *Parabole rectilinea* in HREBLAY & RONKAY (1998: 216). Holotype male, Nepal: “Ganesh Himal, near Godlang” (CGR).
- regina* Hreblay et L. Ronkay, 1999 – Described as *Kisegira regina* in HREBLAY & RONKAY (1999: 507). Holotype male, Thailand: “Thailand, Changwat Nan, 30 km E of Pua” (CMH).
- remota* Hreblay, 1996 – Described as *Mythimna (Morphopoliana) consanguis remota* in HREBLAY (1996b: 136). Holotype male, Australia: “Australien, Queensland, Mackay” (BMNH).
- remota* Hreblay et L. Ronkay, 1998 – Described as *Lithophane remota* in HREBLAY & RONKAY (1998: 181). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 5 km SW of Tayuling” (CMH).
- renimaculata* Hreblay et Legrain, 1996 – Described as *Mythimna (Pseudaletia) renimaculata* in HREBLAY *et al.* (1996: 96). Holotype male, Nepal: “Nepal, Ganesh Himal, 2 km E of Thangjet” (CMH).
- repeteki* L. Ronkay, Varga et Hreblay, 1998 – Described as *Chilodes repeteki* in RONKAY *et al.* (1998: 272). Holotype female, Turkmenistan: “USSR, Turkmenia, Kara-Kum desert, 20 km SW of Repetek” (HNHM).
- replicata* Hreblay et Yoshimatsu, 1999 – Described as *Analetia (Anapoma) replicata* in HREBLAY *et al.* (1999a: 389). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok, 16 km NW of Fang” (CMH).
- reseri* Hreblay et L. Ronkay, 1998 – Described as *Apamea reseri* in HREBLAY & RONKAY (1998: 249). Holotype male, Nepal: “Ganesh Himal, 2 km E of Thangjet” (CMH).
- reshoefii* Hreblay, 1994 – Described as *Orthosia (Orthosia) reshoefii* in HREBLAY & PLANTE (1994: 26). Holotype female, Afghanistan: “Afghanistan, Dare Porandey” (CMH).
- reticulata* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Owadaglea reticulata* in HREBLAY *et al.* (1999b: 53). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- retunda* Hreblay et L. Ronkay, 1998 – Described as *Blepharosis retunda* in HREBLAY *et al.* (1998b: 141). Holotype male, China: “Li-kiang (China), Provinz Nord-Yuennan” (MAK).
- robusta* Hreblay et Kononenko, 1999 – Described as *Diphtherocome robusta* in HREBLAY & KONONENKO (1999: 676). Holotype male, China: “[China, Prov. Yunnan], Likiang, Prov. Nord-Yuennan” (MAK).
- ronkaygabori* Hreblay et L. Ronkay, 1999 – Described as *Dasypolia (Auropolia) ronkaygabori* in HREBLAY & RONKAY (1999: 521). Holotype male, Nepal: “Nepal, Mechi, Taplejung area, 5 km SW Ramche” (HNHM-GR).
- roseago* L. Ronkay, Varga et Hreblay, 1998 – Described as *Gortyna roseago* in RONKAY *et al.* (1998: 267). Holotype male, Turkmenistan: “Turkmenija, Kara-Kala, Ai-Dere” (ZMK).
- rubiculosa* Hreblay et L. Ronkay, 1999 – Described as *Paranyctycia rubiculosa* in HREBLAY & RONKAY (1999: 543). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Inthanon” (CMH).
- rubrifasciata* Hreblay et L. Ronkay, 2000 – Described as *Taivaleria rubrifasciata* in HREBLAY & RONKAY (2000: 20). Holotype male, Taiwan: “Taiwan, Prov. Ilan, Ming Chyr Forest Recreation Area” (CGYF).
- rubrigrappa* Hreblay et L. Ronkay, 1999 – Described as *Harutaographa caerulea rubrigrappa* in HREBLAY & RONKAY (1999: 492). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok, 18 km NW of Fang” (CMH).

- rubrocinerea* Hreblay et L. Ronkay, 1998 – Described as *Perigrappa (Anorthoa) rubrocinerea* in HREBLAY & RONKAY (1998: 153). Holotype female, Nepal: “Nepal, Taplejung area, Mechi, above Hellok” (CGR).
- rufatrox* Hreblay et L. Ronkay, 1995 – Described as *Dasypolia (Dasypolia) rufatrox* in HREBLAY & RONKAY (1995b: 368). Holotype female, Pakistan: “N-Pakistan, 5 km E of Chorit” (CJP).
- rutilitincta* Hreblay et Yoshimatsu, 1996 – Described as *Mythimna (Hyphilare) rutilitincta* in HREBLAY *et al.* (1996: 104). Holotype male, Nepal: “Nepal, Ganesh Himal, 2 km E of Yurekharka” (CMH).
- saba* Hreblay et Plante, 1996 – Described as *Perigrappa (Harutaeographa) saba* in HREBLAY (1996a: 80). Holotype male, Pakistan: “N-Pakistan, 10 km SW of Astor, Rama” (CMH).
- sagitta* Hreblay et L. Ronkay, 1998 – Described as *Hemiexarnis epiphana sagitta* in HREBLAY & RONKAY (1998: 129). Holotype male, Nepal: “Nepal, Annapurna Himal, 5 km SE of Jomsom, Thadung valley” (CGR).
- salaka* Hreblay et L. Ronkay, 1999 – Described as *Meganyctycia salaka* in HREBLAY & RONKAY (1999: 542). Holotype female, Nepal: “West-Nepal, 11 km N of Dailekh” (CMH).
- sanyibaglya* Hreblay et L. Ronkay, 1998 – Described as *Apamea sanyibaglya* in HREBLAY & RONKAY (1998: 244). Holotype male, Nepal: “Ganesh Himal, between Godlang and Nesim” (CSTK).
- satellitita* Hreblay et L. Ronkay, 2000 – Described as *Fabiania satellitita* in HREBLAY & RONKAY (2000: 28). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Sa Pa” (HNHM).
- schaeferi* Hreblay et L. Ronkay, 1998 – Described as *Xestia schaeferi* in HREBLAY *et al.* (1998b: 126). Holotype male, India: “Tsomgo” (MNHU).
- schistochlora* L. Ronkay, Varga et Hreblay, 1998 – Described as *Polymixis schistochlora* in RONKAY *et al.* (1998: 241). Holotype male, Turkmenistan: “USSR, Turkmenia, Kopet-Dagh Mts, 20 km E of Nochur, Karayalchi valley” (HNHM).
- scriptobella* Hreblay, 1996 – Described as *Perigrappa (Perigrappa) scriptobella* in HREBLAY (1996a: 70). Holotype male, Turkmenistan: “Turkmenistan, Kopet-Dagh Mts, 6 km S of Ipaj-Kala” (CMH). (Drawer with paratypes shown in Fig. 6)
- secreta* Hreblay et Plante, 1995 – Described as *Anaplectoides secreta* in HREBLAY & PLANTE (1995a: 136). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, Lamjura Pass” (CMH).
- secreta* Hreblay et L. Ronkay, 1999 – Described as *Diphtherocome secreta* in HREBLAY & RONKAY (1999: 564). Holotype male, Nepal: “Nesukharka, 12 km S Somsang” (HNHM).
- selecta* Hreblay et Plante, 1995 – Described as *Hermonassa selecta* in HREBLAY & PLANTE (1995b: 540). Holotype male, Nepal: “Nepal, Taplejung area, Lal Kharka” (CJP).
- semivirens* Hreblay et L. Ronkay, 1998 – Described as *Euplexidia semivirens* in HREBLAY & RONKAY (1998: 265). Holotype male, Nepal: “Nepal, Annapurna Himal, between Ghorepani and Deorali” (HNHM).
- sericea* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Elwesia tarka sericea* in HREBLAY *et al.* (1999b: 55). Holotype male, Nepal: “West-Nepal, 14 km N of Dailekh” (CMH).
- sericina* L. Ronkay, Varga et Hreblay, 1998 – Described as *Allophyes sericina* in RONKAY *et al.* (1998: 229). Holotype male, Turkmenistan: “USSR, Turkmenia, Kopet-Dagh Mts, 5 km S of Chuli” (HNHM).
- sericina* Hreblay et L. Ronkay, 1999 – Described as *Daseuplexia chloromagna sericina* in HREBLAY & RONKAY (1999: 539). Holotype male, Nepal: “Nepal, Mechi, Taplejung area, Deorali, 5 km NE Suketar” (HNHM-GR).
- serrata* Hreblay et Kononenko, 1999 – Described as *Diphtherocome serrata* in HREBLAY & KONONENKO (1999: 676). Holotype male, China: “[China, Prov. Yunnan], Likiang, Prov. Nord-Yuennan” (MAK).

- sherpa* Hreblay et L. Ronkay, 1998 – Described as *Raddea sherpa* in HREBLAY & RONKAY (1998: 140). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, 14 km E of Lukla” (CMH).
- sherpa* Hreblay et L. Ronkay, 1999 – Described as *Dasypolia sherpa* in HREBLAY & RONKAY (1999: 519). Holotype male, Nepal: “East-Nepal, Deorali Danda, 6 km NW of Yamphudin” (CMH).
- siamensis* Hreblay, 1998 – Described as *Mythimna (Hyphilare) siamensis* in HREBLAY *et al.* (1998a: 390). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok” (CMH).
- siamica* Hreblay et L. Ronkay, 1999 – Described as *Agrochola siamica* in HREBLAY & RONKAY (1999: 528). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Phahompok, 18 km NW of Fang” (CMH).
- siamica* Hreblay et L. Ronkay, 1999 – Described as *Owadaglea elongata siamica* in HREBLAY & RONKAY (1999: 547). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Inthanon” (CMH).
- signa* Hreblay et L. Ronkay, 2000 – Described as *Nyctycia signa* in HREBLAY & RONKAY (2000: 23). Holotype female, Taiwan: “Taiwan, Prov. Nantou, 3 km SW of Tsuifeng” (CMH).
- similissima* Hreblay et Yoshimatsu, 1996 – Described as *Mythimna (Hyphilare) similissima* in HREBLAY *et al.* (1996: 110). Holotype male, Nepal: “Nepal, Ganesh Himal, 2 km E of Thangjet” (CMH).
- simillima* Hreblay et L. Ronkay, 1995 – Described as *Himalistra simillima* in HREBLAY & RONKAY (1995a: 240). Holotype female, Nepal: “Nepal, Ganesh Himal, 1 km SE of Somdang” (HNHM).
- simonyii* Hreblay, 1998 – Described as *Nyctycia simonyii* in KOBAYASHI *et al.* (1998: 260). Holotype male, Taiwan: “Taiwan, Taoyuan, 7 km E of Fuhsing” (CMH).
- singularis* Hreblay et L. Ronkay, 1998 – Described as *Orthosia singularis* in HREBLAY & RONKAY (1998: 157). Holotype male, Nepal: “Nepal, Ganesh Himal, near Nesim” (CGR).
- siva* Hreblay, 1996 – Described as *Perigrapha (Harutaographa) siva* in HREBLAY (1996a: 80). Holotype female, India: “India, Simla” (BMNH).
- sobeyi* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Hermonassa sobeyi* in HREBLAY *et al.* (1999b: 2). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- sokotrensis* Hreblay, 1996 – Described as *Mythimna (Morphopoliana) sokotrensis* in HREBLAY (1996b: 145). Holotype female, Yemen: “Socotra, Hadibo Plain” (BMNH).
- soluta* Hreblay et L. Ronkay, 1997 – Described as *Himalistra soluta* in HREBLAY & RONKAY (1997: 49). Holotype female, Taiwan: “Taiwan, Prov. Nantou, 5 km SW of Tayiling” (CMH).
- sordescens* Hreblay, 1993 – Described as *Orthosia (Monima) sordescens* in HREBLAY (1993: 79). Holotype male, Russia: “USSR, N-Kaukasus, Stavropol” (CMH).
- stegeri* Hreblay et Thöny, 1995 – Described as *Checupa stegeri* in HREBLAY & THÖNY (1995: 24). Holotype male, Taiwan: “Taiwan, Prov. Taoyuan, 14 km E of Fuhshing” (CMH).
- stevenswani* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Actinotia stevenswani* in HREBLAY *et al.* (1999b: 80). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Sa Pa” (HNHM).
- suavina* Hreblay, 1998 – Described as *Mythimna (Mythimna) suavina* in HREBLAY *et al.* (1998a: 384). Holotype male, Myanmar: “Burmah” (BMNH).
- subapicalis* Hreblay et L. Ronkay, 1998 – Described as *Hyalobole subapicalis* in HREBLAY & RONKAY (1998: 238). Holotype male, Nepal: “Nepal, Arun valley, 12 km N of Chitre” (CMH).
- subcyanea* Hreblay et L. Ronkay, 1999 – Described as *Cryphia (Hymenocryphia) subcyanea* in HREBLAY & RONKAY (1999: 572). Holotype male, Nepal: “East-Nepal, Deorali Danda, Anpan” (CMH).

- subforsteri* Hreblay et L. Ronkay, 1998 – Described as *Xestia subforsteri* in HREBLAY *et al.* (1998b: 127). Holotype male, India: “Tsomgo, obere Lage” (MNHU).
- substriata* Hreblay et L. Ronkay, 1997 – Described as *Apamea aquila substriata* in HREBLAY & RONKAY (1997: 59). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 5 km SW of Tayüling” (CMH).
- subtropica* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Hyalobole subtropica* in HREBLAY *et al.* (1999b: 57). Holotype male, Thailand: “Thailand, Changwat Chian Mai, Mt Doi Inthanon, NP” (CMH).
- sugii* Hreblay et L. Ronkay, 1998 – Described as *Dichagyris sugii* in HREBLAY & RONKAY (1998: 128). Holotype male, Nepal: “Nepal, Annapurna Himal, 10 km SE of Jomsom” (CGR).
- szabokyi* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Nyctycia szabokyi* in HREBLAY *et al.* (1999b: 12). Holotype male, Thailand: “Thailand, Changwat Mae Hong Song, 21 km NW of Pai” (CMH).
- szecsenyii* Hreblay et L. Ronkay, 1998 – Described as *Phlogophora szecsenyii* in HREBLAY *et al.* (1998b: 148). Holotype male, Nepal: “East-Nepal, Deorali Danda, Anpan” (CMH).
- tahiricola* L. Ronkay et Hreblay, 1994 – Described as *Himalistra tahiricola* in HREBLAY *et al.* (1994: 255). Holotype male, Turkey: “Turkey, Prov. Agri, 5 km E of Sarican” (CMH).
- taimacula* Hreblay et L. Ronkay, 2000 – Described as *Sphragifera sigillata taimacula* in HREBLAY & RONKAY (2000: 35). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 3 km SW of Tsuifeng” (CMH).
- taitungensis* Hreblay et L. Ronkay, 1997 – Described as *Plataplecta pulverosa taitungensis* in HREBLAY & RONKAY (1997: 22). Holotype male, Taiwan: “Taiwan, Prov. Taitung, 5 km NW of Lirao” (CMH).
- taiwanensis* Hreblay et L. Ronkay, 1998 – Described as *Hyalobole taiwanensis* in HREBLAY & RONKAY (1998: 241). Holotype male, Taiwan: “Taiwan, Hualien Hsien, Houhuansan, Luoying Lodge” (HNHM).
- taiwanicola* Hreblay et L. Ronkay, 2000 – Described as *Xylopolia bella taiwanicola* in HREBLAY & RONKAY (2000: 16). Holotype male, Taiwan: “Taiwan, Prov. Hualien, Taroko N.P., Hsipan” (HNHM).
- tamangi* Hreblay et Plante, 1996 – Described as *Tricheurois tamangi* in HREBLAY & PLANTE (1996: 665). Holotype male, Nepal: “Nepal, Ganesh Himal, Khurpudanda Pass” (CJP).
- tarka* Hreblay et L. Ronkay, 1998 – Described as *Elwesia tarka* in HREBLAY & RONKAY (1998: 233). Holotype female, Nepal: “Ganesh Himal, 2 km W of Gholjong” (CGR).
- tekke* L. Ronkay, Varga et Hreblay, 1998 – Described as *Periphanes delphinii tekke* in RONKAY *et al.* (1998: 219). Holotype male, Turkmenistan: “Turkmenistan, 50 km N of Ashkhabad” (CMH).
- telortoides* Hreblay et L. Ronkay, 1999 – Described as *Agrochola telortoides* in HREBLAY & RONKAY (1999: 525). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- tengeri* Hreblay, 1999 – Described as *Mythimna (Pseudaletia) tengeri* in HREBLAY *et al.* (1999a: 378). Holotype female, Indonesia: “[Indonesien, Java], Tenger” (NNM).
- tertia* Hreblay et L. Ronkay, 1999 – Described as *Daseuplexia tertia* in HREBLAY & RONKAY (1999: 538). Holotype male, Nepal: “East-Nepal, Surke Danda, 2 km NE of Suketar” (CMH).
- thailandica* Hreblay, 1998 – Described as *Mythimna (Morphopoliana) thailandica* in HREBLAY *et al.* (1998a: 410). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, 23 km NW of Sop Kha, 1 km E of Kop Dong” (CMH).

- thailandica* Hreblay et L. Ronkay, 1999 – Described as *Csorbatype thailandica* in HREBLAY & RONKAY (1999: 551). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Inthanon” (CMH).
- thainympha* Hreblay et L. Ronkay, 1999 – Described as *Diphtherocome thainympha* in HREBLAY & RONKAY (1999: 570). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Inthanon” (CMH).
- thaumasia* Hreblay et L. Ronkay, 1998 – Described as “*Nyctycia*” *thaumasia* in HREBLAY & RONKAY (1998: 218). Holotype female, Nepal: “Nepal, Annapurna Himal, vic. of Lumle” [CGR]. Note: In the original description the depositor has not been indicated. The emerging question has been solved via personal communication of Dr László L. Ronkay.
- theobroma* Hreblay et L. Ronkay, 1997 – Described as *Blepharosis theobroma* in HREBLAY & RONKAY (1997: 47). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 3 km SW of Tsuifeng” (CMH).
- thinicola* Hreblay et L. Ronkay, 1998 – Described as *Cryphia thinicola* in HREBLAY & RONKAY (1998: 125). Holotype male, Nepal: “Nepal, Annapurna Himal, 1 km S of Jomson, Thini vilage” (CGR).
- tibetana* Hreblay et L. Ronkay, 1995 – Described as *Estagrotis canescens tibetana* in HREBLAY & RONKAY (1995a: 250). Holotype male, China: “China, Tibet, 50 km N of Nyalam” (HNHM).
- tibetensis* Hreblay, 1998 – Described as *Mythimna (Hyphilare) tibetensis* in HREBLAY *et al.* (1998a: 401). Holotype male, China: “S. Tibet, Chumbi Valley” (BMNH).
- tibori* Hreblay, 1994 – Described as *Egira tibori* in HREBLAY (1994: 247). Holotype male, Turkey: “Turkey, P. Maras. 1 km N of Kahramanmaras” (CMH).
- tibori* Hreblay et L. Ronkay, 1998 – Described as *Estagrotis tibori* in HREBLAY & RONKAY (1998: 197). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 5 km SW of Tayuling” (CMH).
- tingricola* Hreblay et L. Ronkay, 1998 – Described as *Parexarnis tingricola* in HREBLAY *et al.* (1998b: 124). Holotype male, China: “China, Tibet, 20 km E of Old Tingri” (CJP).
- tinsangla* Hreblay et L. Ronkay, 1999 – Described as *Dasyptolia (Dasymixis) tinsangla* in HREBLAY & RONKAY (1999: 520). Holotype male, Nepal: “Nepal, Mt Kalinchok, 1 km SW of Tinsang Pass” (CMH).
- tonkini* Hreblay et L. Ronkay, 1999 – Described as *Euplexidia tonkini* in HREBLAY & RONKAY (1999: 560). Holotype male, Vietnam: “Vietnam, Tonkin, Fan-si-pan” (CMH).
- transformis* Hreblay et L. Ronkay, 1999 – Described as *Harutaographa yangzisherpani transformis* in HREBLAY & RONKAY (1999: 491). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM-GR).
- triangulifera* Hreblay et L. Ronkay, 1998 – Described as *Owadaglaea triangulifera* in HREBLAY & RONKAY (1998: 226). Holotype male, Nepal: “Nepal, Mechi, Taplejung area, above Yamphudin” (CGR).
- tricorna* Hreblay, Legrain et Yoshimatsu, 1998 – Described as *Mythimna (Hyphilare) tricorna* in HREBLAY *et al.* (1998a: 396). Holotype male, Thailand: “Tahiland [sic!], Changwat Nan, 25 km N of Bo Luang” (CMH).
- trifurcata* Hreblay et L. Ronkay, 1999 – Described as *Xestia trifurcata* in HREBLAY & RONKAY (1999: 486). Holotype male, Nepal: “East-Nepal, Kanchenjunga Himal, Rampuk Kharka” (CMH).
- trimaculata* Hreblay et Kononenko, 1999 – Described as *Diphtherocome biformis trimaculata* in HREBLAY & KONONENKO (1999: 680). Holotype male, China: “[China, Prov. Yunnan], Likiang, Prov. Nord-Yuennan” (MAK).
- trimorpha* Hreblay et L. Ronkay, 1997 – Described as *Lithophane trimorpha* in HREBLAY & RONKAY (1997: 42). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 5 km SW of Tayuling” (CMH).

- tristigmata* Hreblay et L. Ronkay, 1998 – Described as *Agrotis tristigmata* in HREBLAY *et al.* (1998b: 116). Holotype male, China: “China, Tibet, 1 km S of Paiku Tso” (CMH).
- trivibia* Hreblay et L. Ronkay, 1998 – Described as *Euxoa trivibia* in HREBLAY *et al.* (1998b: 112). Holotype male, China: “China, Tibet, 50 km N of Nyalam” (CJP).
- tseringi* Hreblay et L. Ronkay, 1995 – Described as *Dasyptolia (Dasyptolia) tseringi* in HREBLAY & RONKAY (1995b: 365). Holotype female, China: “China, Tibet: 4 km S of Karru-Ochen Tso” (CJP).
- tshava* Hreblay et L. Ronkay, 1999 – Described as *Sidemia stoliczcae tshava* in HREBLAY & RONKAY (1999: 557). Holotype male, Nepal: “Nepal, Annapurna Himal, 4 km NE of Tukuhe” (CMH).
- turcomana* L. Ronkay, Varga et Hreblay, 1998 – Described as *Omphalophana turcomana* in RONKAY *et al.* (1998: 227). Holotype male, Turkmenistan: “USSR, Turkmenia, Kopet-Dagh Mts, 5 km NW of Tutlikala” (HNHM).
- turcomania* L. Ronkay, Yela & Hreblay, 2001 – Described as *Leucochlaena (Leucochlaena) muscosa turcomania* in RONKAY *et al.* (2001: 257). Holotype male, Turkmenistan: “Kopet-Dagh Mts, Arap-Ata, 12 km E Kara-Kala” (HNHM).
- turcomanica* L. Ronkay, Varga et Hreblay, 1998 – Described as *Agrochola turcomanica* in RONKAY *et al.* (1998: 258). Holotype male, Turkmenistan: “USSR, Turkmenia, Kopet-Dagh Mts, Sayvan valley, cca 10 km N of Sayvan” (HNHM).
- turpika* Hreblay, 1991 – Described as *Orthosia (Semiophora) askoldensis turpika* in HREBLAY (1991: 197). Holotype male, Mongolia: “Mongolia, Central aimak, Tsagaan Davaa” (CMH).
- unica* Hreblay et L. Ronkay, 1998 – Described as *Himalaea unica* in HREBLAY *et al.* (1998b: 136). Holotype male, China: “China, Tibet, 10 km E of Nyemo” (CMH).
- unicata* Hreblay et L. Ronkay, 1995 – Described as *Dasyptolia (Dasyptolia) unicata* in HREBLAY & RONKAY (1995b: 359). Holotype male, China: “China, Tibet, 1 km S of Nyalam” (CJP).
- unicolor* Hreblay et L. Ronkay, 1998 – Described as *Lithopolia confusa unicolor* in HREBLAY & RONKAY (1998: 162). Holotype male, Nepal: “Nepal, Ganesh Himal, 1 km E of Gadrang” (CMH).
- unicolor* Gyulai, Hreblay et L. Ronkay, 1998 – Described as *Blepharosis bryocharis unicolor* in HREBLAY *et al.* (1998b: 141). Holotype male, China: “China, Quinghai, Anemaquen shan” (CPGY).
- unicuspis* Hreblay et Plante, 1998 – Described as *Athetis unicuspis* in HREBLAY *et al.* (1998b: 152). Holotype male, China: “China, Tibet, 10 km E of Nyemo” (CJP).
- ursina* Hreblay et L. Ronkay, 1995 – Described as *Dasyptolia (Dasyptolia) ursina* in HREBLAY & RONKAY (1995b: 366). Holotype female, Nepal: “Nepal, Ganesh Himal, Khurpudanda Pass, NO slope” (HNHM).
- vanensis* Hreblay et L. Ronkay, 1992 – Described as *Oncocnemis exacta vanensis* in GYULAI *et al.* (1992: 37). Holotype male, Turkey: “Turkey, Prov. Bitlis, 16 km E of Tatvan” (HNHM).
- vargai* Hreblay et L. Ronkay, 1998 – Described as *Discestra vargai* in HREBLAY & RONKAY (1998: 146). Holotype male, Nepal: “Nepal, Annapurna Himal, Mesokantu Pass” (CGR).
- vargai* Hreblay et L. Ronkay, 1998 – Described as *Conisania vargai* in HREBLAY *et al.* (1998b: 131). Holotype male, China: “Gyantse” (MNHU).
- variabilis* Hreblay et Plante, 1995 – Described as *Himalistra variabilis* in HREBLAY *et al.* (1995: 48). Holotype male, China: “China, Tibet, 1 km S of Nyalam” (CMH).
- variegata* Hreblay et L. Ronkay, 1998 – Described as *Hyalobole variegata* in HREBLAY & RONKAY (1998: 237). Holotype female, Nepal: “Ganesh Himal, Gothen village” (HNHM).
- variegata* Hreblay et L. Ronkay, 1998 – Described as *Oroplexia variegata* in HREBLAY & RONKAY (1998: 256). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 5 km SW of Tayuling” (CMH).

- vasarhelyii* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Elwesia diplostigma vasarhelyii* in HREBLAY *et al.* (1999b: 57). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 6 km W Sa Pa” (HNHM).
- vecorina* Hreblay et Plante, 1998 – Described as *Isochlora vecors vecorina* in HREBLAY *et al.* (1998b: 109). Holotype male, China: “China, Tibet, Mayum La (Pass), E side” (CJP).
- vernalis* Hreblay et L. Ronkay, 1998 – Described as *Nyctycia vernalis* in HREBLAY & RONKAY (1998: 201). Holotype female, Nepal: “Ganesh Himal, near Haku” (CGR).
- vindhya* Hreblay et Legrain, 1999 – Described as *Leucania vindhya* in HREBLAY *et al.* (1999a: 385). Holotype male, India: “[India], M S Wyndia, Plateau Sumakna” (IRSNB).
- violacea* Hreblay et L. Ronkay, 1998 – Described as *Xestia violacea* in HREBLAY & RONKAY (1998: 135). Holotype male, Nepal: “Nepal, Solu Khumbu Himal, Lukla” (CMH).
- violascens* Hreblay et L. Ronkay, 1998 – Described as *Lithophane violascens* in HREBLAY & RONKAY (1998: 180). Holotype female, Nepal: “Ganesh Himal, 1 km E of Gadrang” (CMH).
- violascens* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Nyctycia stenoptera violascens* in HREBLAY *et al.* (1999b: 10). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- viridentata* Hreblay et L. Ronkay, 1997 – Described as *Antivaleria viridentata* in HREBLAY & RONKAY (1997: 45). Holotype male, Taiwan: “Taiwan, Prov. Taitung, Hsiangyang” (CGYF).
- viridicincta* Hreblay et L. Ronkay, 1998 – Described as *Daseuplexia viridicincta* in HREBLAY & RONKAY (1998: 210). Holotype male, Nepal: “Nepal, Langtang, between Cholang Pati and Dimsa” (HNHM).
- viridinigra* Hreblay et L. Ronkay, 1997 – Described as *Polymixis viridinigra* in HREBLAY & RONKAY (1997: 46). Holotype male, Taiwan: “Taiwan, Prov. Nantou, 7 km SW of Tayüling” (CMH).
- viridis* Hreblay et L. Ronkay, 1998 – Described as *Pygmeopolia viridis* in HREBLAY & RONKAY (1998: 160). Holotype male, Nepal: “Nepal, Annapurna Himal, 1.5 km SE of Nangethanti” (CGR).
- viridissima* Hreblay et Thöny, 1995 – Described as *Checupa viridissima* in HREBLAY & THÖNY (1995: 27). Holotype male, Indonesia: “Indonesien/S-Sulawesi, Rantepao” (CMH).
- viridissima* Hreblay et L. Ronkay, 1999 – Described as *Diphtherocome viridissima* in HREBLAY & RONKAY (1999: 569). Holotype male, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- viriditincta* Hreblay et L. Ronkay, 1998 – Described as *Nyctycia viriditincta* in HREBLAY & RONKAY (1998: 203). Holotype male, Nepal: “Nepal, Arun valley, 12 km N of Chitre” (CMH).
- vitalina* Hreblay et Thöny, 1995 – Described as *Checupa vitalina* in HREBLAY & THÖNY (1995: 28). Holotype male, Indonesia: “Indonesien/N-Tengg, Timur, Flores, 9 km südl. Ruteng, Golo Luseng” (CMH).
- vuquangconi* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Elwesia vuquangconi* in HREBLAY *et al.* (1999b: 55). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 6 km W Sa Pa” (HNHM).
- warreni* Hreblay et L. Ronkay, 1999 – Described as *Isolasia warreni* in HREBLAY & RONKAY (1999: 540). Holotype male, India: “Sikkim” (BMNH).
- weixleri* Hreblay et L. Ronkay, 1997 – Described as *Meganephria weixleri* in HREBLAY & RONKAY (1997: 38). Holotype male, Taiwan: “Taiwan, Prov. Nantou, Meimu” (CGYF).
- xanthophylla* Hreblay et L. Ronkay, 1998 – Described as *Xanthia xanthophylla* in HREBLAY & RONKAY (1998: 242). Holotype male, Nepal: “Nepal, 4 km SW of Kalinchok peak” (CMH).
- yangzisherpani* Hreblay, 1998 – Described as *Panthea yangzisherpani* in HREBLAY (1998: 8). Holotype male, Nepal: “East Nepal, Deorali Danda, Tseram” (CMH).

- yangzisherpani* Hreblay et L. Ronkay, 1999 – Described as *Harutaeographa yangzisherpani* in HREBLAY & RONKAY (1999: 490). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Inthanon” (CMH).
- yoshimotoi* Hacker et Hreblay, 1996 – Described as *Perigrapha (Harutaeographa) yoshimotoi* in HREBLAY (1996a: 75). Holotype male, India: “Indien, Himachal Pradesh. Rohtang-Paß, ob. Kothi” (CHH).
- yoshimotoi* Hreblay et L. Ronkay, 1998 – Described as *Elwesia sugii yoshimotoi* in HREBLAY & RONKAY (1998: 235). Holotype male, Taiwan: “Taiwan, Prov. Taichung, Anmashan, Hooping” (CGYF).
- yoshimotoi* Hreblay et L. Ronkay, 1998 – Described as *Conisania ronkayi yoshimotoi* in HREBLAY et al. (1998b: 130). Holotype male, China: “China, Tibet, Thung La (Pass) W side” (CMH).
- ypsiloidea* Hreblay, Peregovits et L. Ronkay, 1999 – Described as *Diarsia ypsiloidea* in HREBLAY et al. (1999b: 5). Holotype female, Vietnam: “Vietnam, Prov. Lao Cai, Fan-si-pan Mts, 14 km NW Sa Pa” (HNHM).
- yuennana* Hreblay et L. Ronkay, 1997 – Described as *Allophytes yuennana* in HREBLAY & RONKAY (1997: 40). Holotype male, China: “Likiang (China), Provinz Nord-Yuennan” (MAK).
- zita* Hreblay et L. Ronkay, 1999 – Described as *Agrochola zita* in HREBLAY & RONKAY (1999: 526). Holotype male, Thailand: “Thailand, Changwat Nan, 30 km E of Pua” (CMH).
- zita* Hreblay et L. Ronkay, 1999 – Described as *Nyctycia flavipicta zita* in HREBLAY & RONKAY (1999: 536). Holotype male, Thailand: “Thailand, Changwat Chiang Mai, Mt Doi Inthanon” (CMH).
- zwicki* Hreblay et L. Ronkay, 1998 – Described as *Spinipalpa zwicki* in HREBLAY et al. (1998b: 129). Holotype male, India: “Sikkim, Yayokang” (MNHU).

Species-group names proposed as replacement

- draudtiana* Hreblay et Plante, 1996 – Proposed as *Oroplexia draudtiana* nom. n. replacing *Blepharidia retrahens* Draudt, 1950 in HREBLAY & PLANTE (1996: 665).
- snelleni* Hreblay, 1996 – Proposed as *Mythimna (Morphopoliana) snelleni* nom. n. replacing *Hecatera impura* Snellen, 1886 in HREBLAY (1996b: 143).

Genus-group names

- Auropolia* Hreblay et L. Ronkay, 1999 – Described in HREBLAY & RONKAY (1999: 521). Type species: *Dasytopia ronkaygabori* Hreblay et L. Ronkay, 1999.
- Blepharomima* Hreblay et L. Ronkay, 1998 – Described in HREBLAY & RONKAY (1998: 212). Type species: *Blepharomima euplexina* Hreblay et L. Ronkay, 1998.
- Blepharomorpha* Hreblay et L. Ronkay, 1998 – Described in HREBLAY et al. (1998b: 142). Type species: *Blepharomorpha machka* Hreblay et L. Ronkay, 1998.
- Bryotypella* Hreblay et L. Ronkay, 1998 – Described in HREBLAY & RONKAY (1998: 213). Type species: *Dryobata leucostica* Moore, 1882.
- Charanyctycia* Hreblay et L. Ronkay, 1998 – Described in HREBLAY & RONKAY (1998: 218). Type species: *Charanyctycia laudeti* Hreblay et L. Ronkay, 1998.
- Csorbatype* Hreblay, Peregovits et L. Ronkay, 1999 – Described in HREBLAY et al. (1999b: 61). Type species: “*Mniotype*” *csorbai* Hreblay et L. Ronkay, 1998.
- Daseutype* Hreblay, Peregovits et L. Ronkay, 1999 – Described in HREBLAY et al. (1999b: 36). Type species: *Daseutype porphyrea* Hreblay, Peregovits et L. Ronkay, 1999.

- Dioszeghyana* Hreblay, 1993 – Described in HREBLAY (1993: 80). Type species: *Monima schmidtii* Diószeghy, 1935.
- Fabiania* Hreblay et L. Ronkay, 2000 – Described in HREBLAY & RONKAY (2000: 24). Type species: *Fabiania pulla* Hreblay et L. Ronkay, 2000.
- Furcochlaena* L. Ronkay, Yela & Hreblay, 2001 – Described in RONKAY *et al.* (2001: 258). Type species: *Heliophobus fallax* Staudinger, 1870.
- Hiemopolia* Hreblay et L. Ronkay, 1999 – Described in HREBLAY & RONKAY (1999: 510). Type species: *Lithopolia costimacula* Yoshimoto, 1994.
- Himalaea* Hreblay et L. Ronkay, 1998 – Described in HREBLAY *et al.* (1998b: 135). Type species: *Himalaea unica* Hreblay et L. Ronkay, 1998.
- Kisegira* Hreblay et L. Ronkay, 1999 – Described in HREBLAY & RONKAY (1999: 504). Type species: *Kisegira penicillata* Hreblay et L. Ronkay, 1999.
- Loxopamea* Hreblay et Plante, 1995 – Described in HREBLAY & PLANTE (1995b: 544). Type species: *Loxopamea albitractora* Hreblay et Plante, 1995.
- Meganyctycia* Hreblay et L. Ronkay, 1998 – Described in HREBLAY & RONKAY (1998: 207). Type species: *Meganyctycia forcipata* sp. n.
- Morphopholiana* Hreblay et Legrain, 1996 – Described in HREBLAY (1996b: 134). Type species: *Hadena consanguis* Guenée, 1852.
- Opacographa* Hreblay, 1996 – Described in HREBLAY (1996a: 72). Type species: *P. cilissa* Püngeler, 1917.
- Pantheana* Hreblay, 1998 – Described in HREBLAY (1998: 7). Type species: *Pantheana yangzisherpani* Hreblay, 1998. Remark: In the heading of the description the type species is incorrectly referred to as “*Panthea yangzisherpani* sp. n.” but this is clearly an error as all other mentions in the article (at least seven times) use *Pantheana* indicating the correct combination. The combination as written out on top of page 8 is considered a lapsus, and it neither affects the availability of the genus-group name nor the original (intended) combination of the species-group name. The name *Pantheana* is available for nomenclatural purposes under the authorship of Hreblay, 1988 and the type species of the genus is *Pantheana yangzisherpani*.
- Parabole* Hreblay et L. Ronkay, 1998 – Described in HREBLAY & RONKAY (1998: 215). Type species: *Parabole rectilinea* Hreblay et L. Ronkay, 1998.
- Paranyctycia* Hreblay et L. Ronkay, 1998 – Described in HREBLAY & RONKAY (1998: 214). Type species: *Paranyctycia orbiculosa* Hreblay et L. Ronkay, 1998.
- Pectinipolia* Hreblay et L. Ronkay, 1999 – Described in HREBLAY & RONKAY (1999: 510). Type species: *Lithopolia indistincta* Hreblay et L. Ronkay, 1998.
- Peperina* Hreblay, 1992 – Described in HREBLAY (1992b: 542). Type species: *Cerastis torrida* Lederer, 1857.
- Potnyctycia* Hreblay et L. Ronkay, 1998 – Described in HREBLAY & RONKAY (1998: 204). Type species: *Potnyctycia porphyrea* Hreblay et L. Ronkay, 1998.
- Prodigithymna* Hacker, L. Ronkay et Hreblay, 2002 – Described in HACKER *et al.* (2002: 185). Type species: *Leucania prominens* Walker, 1856.
- Pygmeopolia* Hreblay et L. Ronkay, 1998 – Described in HREBLAY & RONKAY (1998: 159). Type species: *Pygmeopolia viridis* Hreblay et L. Ronkay, 1998.
- Taivaleria* Hreblay et L. Ronkay, 2000 – Described in HREBLAY & RONKAY (2000: 19). Type species: *Taivaleria rubrifasciata* Hreblay et L. Ronkay, 2000.
- Thaipolia* Hreblay et L. Ronkay, 1999 – Described in HREBLAY & RONKAY (1999: 510). Type species: *Lithopolia inversa* Hreblay et L. Ronkay, 1999.
- Transeuplexia* Hreblay et Plante, 1995 – Described in HREBLAY & PLANTE (1995a: 139). Type species: no type species. Remark: In the original description “*Transeuplexia violascens* (Boursin,

1964) comb. n.” and “*Transeuplexia grisea* n. sp.” are mentioned, but no type species was selected. Thus, according to the ICZN (1999), article 13.3., the genus-group name *Transeuplexia* under the authorship of Hreblay et Plante is not available for any nomenclatural purpose.

DISCUSSION

Primary types according to their geographical origin

From the localities of his primary types it is obvious that the exploratory activity of M. Hreblay was focused on the Asian fauna, especially the region of the Himalayas and some adjacent territories (Table 1). By far the highest number of taxa was described from Nepal (235; 39%), the country where he displayed great activity organising various expeditions and concentrated on the exploration of the Noctuidae fauna. Almost equivalent effort has been made in neighbouring territories: China (Tibet) (76), Taiwan (62) and Thailand (53) (altogether 191; 32%) to where his interest turned after the great scientific and financial success of the Nepalese expeditions. Practically all the specimens that became primary types and deposited in his own collection were collected by him or his staff (Fig. 7).

Other countries as Pakistan (18) and Turkmenistan (28) with notable number of taxa described by M. Hreblay refer to the previous series of expedi-



Fig. 7. The staff of the Nepalese expedition in March, 1995, at Nangethanthi, 2445 m, Annapurna Region; Dr Hreblay (indicated by arrow) amongst the sherpa porters (photo: L. Németh)

tions by Hungarian entomologists, who (after several successful trips in the Balkans, Mongolia, Turkey and the Caucasus region) also started collecting insects in Central Asia. M. Hreblay participated in several of these trips. The remaining countries with comparatively low number of described taxa testify that M. Hreblay had several colleagues and collaborators who supplied material from areas he did not visit himself or he was working on specimens originating from various museum collections as an expert of noctuid taxonomy. The exception is Vietnam from where also a high number of taxa has been described (54; ~10%). In this country staff members of the HNHM have carried out many expeditions, and M. Hreblay was also involved in the elaboration of the Vietnamese Noctuidae fauna.

Having databased the collection it is evident that there are still hundreds of specimens with great taxonomic significance. Undescribed taxa (already separated by M. Hreblay himself) also await formal description and to be named, not only from the Himalayas but also from subtropical and tropical regions of Indonesia, Thailand, the Philippines, and from the Tibetan Plateau. This highly interesting material can be accessed and studied by scientists when the catalogue of the entire Hreblay Collection will have been published.

Table 1. Countries with type localities of taxa described by Dr Hreblay (with the indication of the existing names' number).

Afghanistan (2)	Malaysia (1)	Taiwan (62)
Australia (2)	Mauritius (1)	Tajikistan (3)
China (76)	Mongolia (5)	Tanzania (1)
Ethiopia (1)	Morocco (1)	Thailand (53)
Fiji (1)	Myanmar (1)	Turkey (7)
France (1)	Nepal (235)	Turkmenistan (28)
Greece (2)	Pakistan (18)	Uzbekistan (1)
India (24)	Papua New Guinea (1)	Vietnam (54)
Indonesia (11)	Philippines (6)	Yemen (1)
Laos (1)	Russia (2)	

Primary types according to their depositories

Márton Hreblay ran a successful private enterprise for collecting, databasing, setting, and preparing natural history material, especially invertebrates. He graduated as an agricultural engineer and therefore his interest turned to Lepidoptera, as they are also important for agricultural management. As his own knowledge has grown in identifying the members of the Palaearctic and Oriental fauna his reputation has also grown and thus became an acknowledged specialist of the cosmopolitan noctuid genus *Mythimna* Ochsenheimer, 1819 and other closely allied genera (Noctuidae: Hadeninae). Therefore he created a large pri-

vate reference collection of his beloved noctuid moths from temperate and subtropical Asia. However, based on the material left behind he wanted to have a collection worldwide in scope.

During his scientific activity he was a fervent collector thus not surprisingly most new taxa he published were from his own collection (Table 2). Therefore almost one half of the name bearing types of taxa he described became deposited in the Hreblay collection (278; 46%). Another large share was of the Lepidoptera collection of the HNHM (118; 19.6%) indicating that the material he studied for describing new taxa originated not from expeditions he organised, but the activities of the museum's staff. M. Hreblay was also in contact with Gábor Ronkay (the younger brother of the eminent noctuid specialist László Ronkay), who ran an enterprise similar to that of M. Hreblay. Via the scientific co-operation of M. Hreblay and László Ronkay a significant number of primary types has been deposited in the Gábor Ronkay Collection (67; 11%). We mention here that one of his sponsors, Jacques Plante (1920–2003) also had a working relationship with M. Hreblay on noctuid faunistics and taxonomy. They pioneered together mainly the fauna of Pakistan and the Himalayas. Their joint efforts at the time also resulted in a number of primary types landing in the private collection of the sponsor (36; 6%). After the tragical death of M. Hreblay, Mr. Plante could safely arrange the future of his collection towards the end of his life as it was acquired for a public museum thanks to authorities of the City of Geneva (LANDRY 2001). Resulting from this event and the most recent placing of the Hreblay collection in the HNHM it can be remarked that by far the largest part of M. Hreblay's primary types became deposited in public institutions (510; 85%), and only a smaller fragment remained in private collections (92; 15%).

Table 2. Depositories of primary type specimens of taxa described by M. Hreblay with the number of their holdings. Abbreviations are keyed in the main text.

BMNH (24)	CSTK (1)	NIAES (2)
CCN (1)	CWS (1)	NMNS (1)
CGR (68)	HNHM (111)	NNM (2)
CGYF (18)	HNHM-GR (6)	NSMT (2)
CHH (4)	HNHM-PGY (2)	UO (1)
CJP (36)	IRSNB (1)	ZIN (1)
CMF (2)	MAK (17)	ZMH (2)
CMH (278)	MNHU (12)	ZMK (1)
CPGY (3)	NHMW (1)	ZSM (4)

*

Acknowledgements – The authors express their sincere thanks to the heirs of the late Márton Hreblay, especially the widow Mrs Judit Heck (Érd, Hungary) for her permission to study the col-

lection, and Mr Lajos Németh (Zalaszentmihály, Hungary) for comments and Figure 7. We are also grateful to our colleagues for providing illustrations (Ms Magdolna Katus and Mr Zoltán György) and commenting the manuscript (Dr György Makranczy and Dr László Ronkay).

REFERENCES

- BÁLINT ZS., KATONA G. & RONKAY L. 2013: The scientific publications of dr Márton Hreblay (1963–2000) on Lepidoptera. – *Esperiana Buchreihe zur Entomologie* **18**: 270–273.
- GYULAI P. & †HREBLAY M. 2001: Description of two new species of the genus *Trichanarta* Hampson, 1896 (Lepidoptera: Noctuidae, Hadeninae). – *Esperiana Buchreihe zur Entomologie* **8**: 709–714.
- GYULAI P., HREBLAY M. & RONKAY L. 1992: New taxa and synonyms in the *Oncocnemis exacta* Christoph, 1887 group (Lepidoptera, Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* **38**(1–2): 33–46.
- HACKER H., RONKAY L. & †HREBLAY M. 2002: *Noctuidae Europaeae volume 4 Hadeninae I.* – Entomological Press, Sorø, 419 pp.
- HREBLAY M. 1990: *Mythimna* (*Aletia*) *jutka* sp. n. aus der Mongolei (Lepidoptera, Noctuidae: Hadeninae). – *Annales historico-naturales Musei nationalis Hungarici* **81**: 125–131.
- HREBLAY M. 1991: Neue Taxa aus der Gattung *Orthosia* Ochsenheimer, 1816 (s. I.) (Lepidoptera, Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* **37**(3–4): 193–203.
- HREBLAY M. 1992a: Paläarktische Taxa der *Mythimna* (*Aletia*) *pallens* L. – *impura* Hb. – Artengruppe (Lepidoptera: Noctuidae). – *Esperiana Buchreihe zur Entomologie* **3**: 513–529.
- HREBLAY M. 1992b: Neue Taxa und Synonyme der Gattung *Conistra* Hübner, [1821] (Lepidoptera: Noctuidae). – *Esperiana Buchreihe zur Entomologie* **3**: 531–544.
- HREBLAY M. 1993: Neue Taxa aus der Gattung *Orthosia* Ochsenheimer, 1816 (s. I.) II. (Lepidoptera, Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* **39**(1–4): 71–90.
- HREBLAY M. 1994: New taxa of the tribe *Orthosiini*, IV. (Lepidoptera, Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* **40**(3): 241–252.
- HREBLAY M. 1996a: Neue Paläarktische Taxa aus der Gattung *Perigrapha* Lederer, 1857 (Lepidoptera, Noctuidae). – *Esperiana Buchreihe zur Entomologie* **4**: 65–94.
- HREBLAY M. 1996b: Revision der *Mythimna consanguis*-, *languida*-, *madensis*-, *natalensis*-Artengruppe (*Morphopoliana* subgen. n.) (Lepidoptera, Noctuidae). – *Esperiana Buchreihe zur Entomologie* **4**: 133–158.
- HREBLAY M. 1996c: On the taxonomy of the *Haderonia culta* (Moore, 1881) species group (Lepidoptera, Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* **42**(1): 65–72.
- HREBLAY M. 1998: Eine neue Gattung und Art aus Nepal. – *Facetta* **9**(2): 7–11.
- HREBLAY M. & KONONENKO V. S. 1995: A new species from the genus *Perigrapha* Lederer, 1857 (Lepidoptera, Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* **41**(2): 115–118.
- HREBLAY M. & KONONENKO V. S. 1999: The Chinese species of the *Diphtherocome pallida* and *D. discibrunnea* species-groups (Lepidoptera, Noctuidae, Acronictinae) in the collection of the Zoologisches Forschungsinstitut und Museum Alexander Koenig, Bonn. – *Esperiana Buchreihe zur Entomologie* **7**: 669–686.
- HREBLAY M., LEGRAIN A. & YOSHIMATSU S.-I. 1996: Beschreibung von vierzehn neuen Arten mit Übersicht der Artengruppen aus dem Komplex *Mythimna* Ochsenheimer, 1816. Festlegungen vierzehn Lectotypen, neue Synonymen und Kombinatoren aus der Himalaya Region. – *Annales historico-naturales Musei nationalis Hungarici* **88**: 89–126.
- HREBLAY M., LEGRAIN A. & YOSHIMATSU S.-I. 1998a: Beschreibung von achtundzwanzig neuen Arten und einer Unterart mit Übersicht der Artengruppen aus dem *Mythimna* Ochsenheimer,

- 1816-Komplex aus der südostasiatischen Region. Festlegung von vierzehn Lectotypen, zwei Neotypen, neuen Synonymen und Kombinationen. (I.) – *Esperiana Buchreihe zur Entomologie* 6: 381–432.
- HREBLAY M., LEGRAIN A. & YOSHIMATSU S.-I. 1999a: Beschreibung von zwölf neuen Arten und zwei Unterarten mit einer Übersicht über die Artengruppen des Mythimna Ochsenheimer, 1816-Komplexes der südostasiatischen Region (II.) (Lepidoptera: Noctuidae). – *Esperiana Buchreihe zur Entomologie* 7: 377–398.
- HREBLAY M., PEREGOVITS L. & RONKAY L. 1999b: New genera and species of Noctuidae from Vietnam, Thailand and Nepal (Lepidoptera). – *Acta Zoologica Academiae Scientiarum Hungaricae* 45(1): 1–96.
- HREBLAY M. & PLANTE J. 1994: New taxa of the genus *Orthosia* Ochsenheimer, 1816 (s.1.) III. (Lepidoptera, Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* 40(1): 21–27.
- HREBLAY M. & PLANTE J. 1995a: Description de sept espèces nouvelles du Népal et d'un genre nouveau (Lépidoptères, Noctuidae). – *Lambillionea* 95: 135–144.
- HREBLAY M. & PLANTE J. 1995b: Description de 12 espèces nouvelles du Népal et de Chine et d'un genre nouveau (Lépidoptères, Noctuidae). – *Lambillionea* 95: 539–550.
- HREBLAY M. & PLANTE J. 1996: Description de huit espèces et d'une sous-espèce nouvelles de la région himalayenne; désignation de cinq nouveaux synonymes et de trois lectotypes (Lépidoptères, Noctuidae). – *Lambillionea* 96(4): 662–675.
- HREBLAY M., PLANTE J. & RONKAY L. 1994: New taxa of the genus *Himalistra* Hacker & Ronkay, 1993 (Lepidoptera, Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* 40(3): 253–263.
- HREBLAY M., PLANTE J. & RONKAY L. 1995: New taxa of the genus *Himalistra* Hacker & Ronkay, 1993 (Lepidoptera, Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* 41(1): 47–62.
- HREBLAY M. & RONKAY L. 1995a: On the taxonomy of the genera *Himalistra* Hacker & Ronkay, 1993, and *Etagrotis* Nye, 1975 (Lepidoptera, Noctuidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* 41(3): 235–250.
- HREBLAY M. & RONKAY L. 1995b: New species of *Dasypolia* Guenée, 1852 (s.l.) from the Himalayan region (Lepidoptera, Noctuidae, Xylenini). – *Acta Zoologica Academiae Scientiarum Hungaricae* 41(4): 349–378.
- HREBLAY M. & RONKAY L. 1997: New Noctuidae (Lepidoptera) species from Taiwan and the adjacent areas. – *Acta Zoologica Academiae Scientiarum Hungaricae* 43(1): 21–83.
- HREBLAY M. & RONKAY L. 1998: Noctuidae from Nepal. Moths of Nepal, Volume 5 – *Tinea* 15 (Supplement 1): 117–314.
- HREBLAY M. & RONKAY L. 1999: Neue trífide Noctuidae aus dem himalayischen Raum und der südostasiatischen Region (Lepidoptera: Noctuidae). – *Esperiana Buchreihe zur Entomologie* 7: 485–620.
- HREBLAY M. & RONKAY L. 2000: New Noctuidae species and subspecies from Taiwan and the adjacent areas II (Lepidoptera). – *Insecta Koreana* 17(1/2): 1–38.
- HREBLAY M., RONKAY L. & PLANTE J. 1998b: Contribution to the Noctuidae (Lepidoptera) fauna of Tibet and the adjacent regions. (II.) A systematic survey of the Tibetan Noctuidae fauna based on the material of the Schäfer-expedition (1938–1939) and recent expeditions (1993–1997) (Lepidoptera: Noctuidae). – *Esperiana Buchreihe zur Entomologie* 6: 69–184.
- HREBLAY M. & THÖNY H. 1995: Sechs neue Arten von *Checupa* Moore 1867 aus Südost-Asien (Lepidoptera, Noctuidae). – *Facetta* 6(2): 21–32.
- ICZN [= International Commission on Zoological Nomenclature] 1999: *International Code of Zoological Nomenclature*. – La Garangola, Padova, 126 pp.

- KOBAYASHI H., OWADA M. & HREBLAY M. 1998: On the genus *Nyctycia* Hampson, 1906 (Lepidoptera, Noctuidae, Cuculiinae) in Taiwan, with descriptions of two new species and three new subspecies. – *Tinea* 15(3): 256–265.
- LANDRY B. 2001: Jacques Plante collection to the Muséum d'histoire naturelle, Geneva, Switzerland. – *News of the Lepidopterists' Society* 42(2): 53.
- RONKAY L., YELA J. L. & HREBLAY M. † 2001: *Noctuidae Europaeae. Volume 5 Hadeninae II*. – Entomological Press, Sorø, 452 pp.
- RONKAY L., VARGA Z. & HREBLAY M. 1998: Twenty two new species and six new subspecies of Noctuidae from Turkmenistan and adjacent regions (Lepidoptera). – *Acta Zoologica Academiae Scientiarum Hungaricae* 44(3): 205–281.
- YOSHIMATSU S.-I. & HREBLAY M. 1996a: A new species of the genus *Mythimna* Ochsenheimer (Lepidoptera, Noctuidae) from Thailand. – *Transactions of the Lepidopterological Society, Japan* 47(1): 13–16.
- YOSHIMATSU S.-I. & HREBLAY M. 1996b: Further two species of the genus *Mythimna* (Lepidoptera, Noctuidae) from Taiwan. – *Tinea* 14(4): 264–267.
- YOSHIMATSU S.-I. & HREBLAY M. 1998: Three new species of the genus *Mythimna* (Lepidoptera, Noctuidae) from China. – *Transactions of the Lepidopterological Society, Japan* 49(1): 9–15.